



# Geochemistry, tectonic setting, and petrogenesis of the Coldbrook Group and associated plutons, southern New Brunswick, Canada: A unique Late Ediacaran magmatic episode in Avalonia

## Géochimie, cadre tectonique et pétrogenèse du Groupe de Coldbrook et des plutons associés, sud du Nouveau-Brunswick, Canada : un épisode magmatique unique de l'Édiacarien supérieur en Avalonie

Alicia P. Escribano and Sandra M. Barr

Volume 59, 2023

URI: <https://id.erudit.org/iderudit/1108392ar>  
DOI: <https://doi.org/10.4138/atlgeo.2023.010>

[See table of contents](#)

Publisher(s)

Atlantic Geoscience Society

ISSN

2564-2987 (digital)

[Explore this journal](#)

Cite this article

Escribano, A. & Barr, S. (2023). Geochemistry, tectonic setting, and petrogenesis of the Coldbrook Group and associated plutons, southern New Brunswick, Canada: A unique Late Ediacaran magmatic episode in Avalonia. *Atlantic Geoscience*, 59, 241–257. <https://doi.org/10.4138/atlgeo.2023.010>

Article abstract

The Coldbrook Group is a unique suite of late Ediacaran volcanic and epiclastic rocks deposited in an extensional setting in the Avalonian Caledonia terrane of southern New Brunswick. It is informally divided into a lower part composed of mainly andesitic to dacitic tuffs and epiclastic sedimentary rocks intruded by bimodal plutons, and an upper part composed mainly of mafic flows and tuffs interlayered with but mainly overlain by rhyolitic flows and tuffs and minor epiclastic sedimentary rocks. Recent U–Pb (zircon) dating of four formations in the Coldbrook Group and a subvolcanic granitic dome led to the proposal that the lower and most of the upper Coldbrook Group and associated plutons were formed within <760 kyr at about 551.5 Ma whereas the uppermost rhyolite (Fundy Trail Parkway rhyolite) and underlying basalt of the Hosford Brook Formation formed in a younger event at about 549.5 Ma. Overlapping dates and chemical data also suggest that the subvolcanic granitic domes are the plutonic counterparts of the ca. 551.5 Ma felsic volcanic rocks whereas the other granitic plutons represent extracted melts that evolved separately. Whole-rock geochemical and isotopic data combined with previous field work and map information and integrated with the geochronological data provide new insights into the genesis of these magmas and their change from intermediate to bimodal magmatism. The data indicate that the volcanic and plutonic rocks are mainly tholeiitic and were derived from varying proportions of juvenile melts and recycled older arc lithosphere. The mafic rocks have chemical similarities to flood basalts which inherited some calc-alkalic signatures from older arc-like rocks. Intermediate magmas that formed much of the lower Coldbrook Group had larger lithospheric inputs and thus have mainly calc-alkalic signatures. The Vernon Mountain, Blackall Lake and Silver Hill area rhyolites represent evolved melts derived from the lower Coldbrook Group magmas, and their chemical differences are interpreted to result from melt extraction and solid-liquid dripping magmatic processes. Enhanced extension and thinning led to formation of less lithosphere-contaminated melts that formed the ca. 549.5 Ma bimodal units of the upper Coldbrook Group.

# Geochemistry, tectonic setting, and petrogenesis of the Coldbrook Group and associated plutons, southern New Brunswick, Canada: a unique Late Ediacaran magmatic episode in Avalonia

ALICIA P. ESCRIBANO<sup>1,2\*</sup> AND SANDRA M. BARR<sup>1</sup>

1. Department of Earth and Environmental Science, Acadia University; Wolfville, Nova Scotia B4P 2R6, Canada

2. Present address: Department of Earth Sciences, Memorial University, St. John's, Newfoundland and Labrador A1B 3X7, Canada

\*Corresponding author <adpescribano@mun.ca>

*Date received: 28 August 2023* † *Date accepted: 19 October 2023*

## ABSTRACT

The Coldbrook Group is a unique suite of late Ediacaran volcanic and epiclastic rocks deposited in an extensional setting in the Avalonian Caledonia terrane of southern New Brunswick. It is informally divided into a lower part composed of mainly andesitic to dacitic tuffs and epiclastic sedimentary rocks intruded by bimodal plutons, and an upper part composed mainly of mafic flows and tuffs interlayered with but mainly overlain by rhyolitic flows and tuffs and minor epiclastic sedimentary rocks. Recent U–Pb (zircon) dating of four formations in the Coldbrook Group and a subvolcanic granitic dome led to the proposal that the lower and most of the upper Coldbrook Group and associated plutons were formed within <760 kyr at about 551.5 Ma whereas the uppermost rhyolite (Fundy Trail Parkway rhyolite) and underlying basalt of the Hosford Brook Formation formed in a younger event at about 549.5 Ma. Overlapping dates and chemical data also suggest that the subvolcanic granitic domes are the plutonic counterparts of the ca. 551.5 Ma felsic volcanic rocks whereas the other granitic plutons represent extracted melts that evolved separately. Whole-rock geochemical and isotopic data combined with previous field work and map information and integrated with the geochronological data provide new insights into the genesis of these magmas and their change from intermediate to bimodal magmatism. The data indicate that the volcanic and plutonic rocks are mainly tholeiitic and were derived from varying proportions of juvenile melts and recycled older arc lithosphere. The mafic rocks have chemical similarities to flood basalts which inherited some calc-alkalic signatures from older arc-like rocks. Intermediate magmas that formed much of the lower Coldbrook Group had larger lithospheric inputs and thus have mainly calc-alkalic signatures. The Vernon Mountain, Blackall Lake and Silver Hill area rhyolites represent evolved melts derived from the lower Coldbrook Group magmas, and their chemical differences are interpreted to result from melt extraction and solid-liquid dripping magmatic processes. Enhanced extension and thinning led to formation of less lithosphere-contaminated melts that formed the ca. 549.5 Ma bimodal units of the upper Coldbrook Group.

## RÉSUMÉ


Le groupe de Coldbrook est une succession de roches volcaniques et épicastiques de l'Édiacarien tardif qui se sont déposées à l'intérieur d'un milieu de distension dans le terrane avalonien de Caledonia dans le sud du Nouveau-Brunswick. Le groupe est officiellement divisé en une section inférieure composée de tufs principalement andésitiques à dacitiques et de roches sédimentaires épicastiques pénétrés par des plutons bimodaux, et en une section supérieure essentiellement constituée de tufs et de coulées mafiques interstratifiées avec des tufs et des coulées rhyolitiques ainsi qu'une quantité modeste de roches sédimentaires épicastiques qui, toutefois, les recouvre principalement. Une datation U–Pb (sur zircon) des quatre formations dans le groupe de Coldbrook et d'un pluton granitique subvolcanique a abouti à la proposition que la section inférieure et la majeure partie de la section supérieure du groupe de Coldbrook et les plutons associés se sont formés il y a moins de 760 milliers d'années, soit il y a environ 551,5 Ma, alors que la rhyolite sommitale (rhyolite de la Promenade du Sentier Fundy) et le basalte sous-jacent de la Formation de Hosford Brook se sont formés au cours d'un épisode plus récent il y a environ 549,5 Ma. Les dates qui se chevauchent et les données chimiques permettent par ailleurs de supposer que les intrusions granitiques subvolcaniques constituent les équivalents plutoniques des roches volcanofelsiques d'il y a environ 551,5 Ma tandis que les plutons granitiques représentent des magmas extraits ayant évolué séparément. Des données isotopiques et géochimiques de roche totale combinées à de l'information cartographique et de l'information émanant de travaux sur le terrain antérieurs intégrées aux données chronologiques livrent de nouveaux renseignements sur la genèse de ces magmas et leur transformation d'un magmatisme intermédiaire à bimodal. Les données révèlent que les roches volcaniques et plutoniques sont

principalement tholéïtiques et qu'elles sont provenues de proportions variables de magmas juvéniles et de la lithosphère d'un arc âgé recyclé. Les roches mafiques ont des similarités chimiques avec les basaltes de plateau et les signatures calco-alkalines héritées des roches d'arc volcanique âgées. Les magmas intermédiaires qui constituent la section inférieure du groupe de Coldbrook ont bénéficié d'afflux lithosphériques plus substantiels et affichent en conséquence des signatures essentiellement calco-alkalines. Les rhyolites des secteurs du mont Vernon, du lac Blackall et de la colline Silver représentent des magmas provenant de magmas de la section inférieure du groupe de Coldbrook et leurs différences chimiques sont interprétées comme un résultat des processus magmatiques de stillation solide-liquide et d'extraction de magma. L'extension et l'amincissement accrus ont entraîné la formation de magmas moins contaminés par la lithosphère qui ont créé les unités bimodales il y a environ 549,5 Ma dans la section supérieure du groupe de Coldbrook.

## INTRODUCTION

Volcanic and epiclastic rocks of the Coldbrook Group and associated plutons underlie about half of the Avalonian Caledonia terrane of southern New Brunswick (Fig. 1). These voluminous volcanic and plutonic rocks formed in a large intraplate magmatic event that accompanied extension

and the opening of sedimentary basins throughout Avalonia in the northern Appalachian orogen during the late Ediacaran (Barr and White 1999a; van Staal *et al.* 2021; Beranek *et al.* 2023; Alvaro *et al.* 2023; Escribano *et al.* 2023). However, elsewhere, in contrast to the Caledonia terrane, only comparatively minor magmatic activity occurred during the late Ediacaran. Also, in other parts of Avalonia,



**Figure 1.** Simplified geological map of the Caledonian Highlands showing the Coldbrook Group and related plutons, younger rocks, and older arc rocks of the Broad River and Lumsden groups after Barr and White (1999a, b, 2004), Barr *et al.* (2020), and Park *et al.* (2014), including some regional structural features. Locations of precisely dated samples from Escribano *et al.* (2023) are shown: 1,  $551.57 \pm 0.23$  Ma (sample NB21-451); 2,  $551.38 \pm 0.24$  Ma (sample NB18-381); 3,  $551.71 \pm 0.19$  Ma (sample NB20-438); 4,  $551.70 \pm 0.20$  Ma (sample NB20-426); 5,  $549.18 \pm 0.09$  Ma (sample NB19-390). Samples labelled 2 and 5, as well as 6 (NB 15-350) and 7 (NB19-391) also have new Sm-Nd isotopic data reported in Table 1. Abbreviation (on inset): CCFS, Cobequid-Chedabucto Fault system.

arc-related magmatism was recorded between ca. 620 Ma and 570 Ma (“main arc phase” of van Staal *et al.* 2021), whereas in the Caledonia terrane, arc magmatism appears to have ended by ~615 Ma and was followed by a magmatic gap until ~551.5 Ma (Escribano *et al.* 2023). Although previous less precise U–Pb zircon dating had indicated that the Coldbrook Group and related plutons formed over a wider late Ediacaran age range of ~560 to 550 Ma (Bevier and Barr 1990; Barr *et al.* 1994, 2019, 2020; Miller *et al.* 2000), Escribano *et al.* (2023) showed that four volcanic samples from throughout the stratigraphic succession (lower Coldbrook Group and Silver Hill Formation and from a subvolcanic intrusion (dome) crystallized in

<760 kyr at ca. 551.5 Ma, and was followed by a large bimodal volcanic event represented by the Hosford Brook Formation and the uppermost rhyolite in the Silver Hill Formation, here termed the Fundy Trail Parkway rhyolite, at ca. 549.5 Ma (Fig. 2).

Previous geochemical studies of the Coldbrook Group and related plutons showed ambiguous within-plate to arc-related signatures and did not clearly identify their petrogenesis and tectonic setting (Currie and Eby 1990; Barr and White 1999a). However, the magmatic rocks were inferred to have formed in a within-plate extensional environment consistent with rift-related sedimentation and minor magmatism recorded elsewhere in Avalonia;

UPPER COLDBROOK GROUP		Previously published	Escribano <i>et al.</i> (2023)
FTPr	<b>Silver Hill Formation</b> ( <i>Fundy Trail Parkway rhyolite</i> ): pink to grey flow-banded rhyolite	548 ± 1 (1)	549.18 ± 0.09 (NB19-390)
HBmv	<b>Hosford Brook Formation</b> : amygdaloidal to massive basalt, with less abundant mafic tuff		
SHfv	<b>Silver Hill Fm</b> ( <i>Silver Hill, Vernon Mountain, &amp; Blackall Lake areas</i> ): pink to red to grey, commonly flow-banded rhyolite and rhyolitic tuff; minor grey to pale brown laminated siltstone and chert	559 ± 5 (2) 555 ± 3 (4) 556 ± 2 (5)	551.70 ± 0.20 (NB20-426)
BUvs	<b>Burley Lake Formation</b> : amygdaloidal basalt flows; locally interlayered rhyolite and red conglomerate and sandstone		
DBv	<b>Dolan Brook Fm</b> : interlayered amygdaloidal basalt, basaltic tuff, rhyolite, & rhyolitic to dacitic tuff	553.3 ± 3 (5) 552.8 ± 1.6 (5)	
BWmv	<b>Browns Lake Formation</b> : green epidotized basalt and andesite; locally includes plagioclase porphyry and green dacitic sheets		
	UPHAM MOUNTAIN PLUTON (mainly granite)	551 ± 5 (2)	
	BONNELL BROOK PLUTON (Mainly granite, minor diorite & gabbro)	550 ± 1 (1) 552.5 ± 1.5 (5)	551.71 ± 0.19 (NB20-438)
	MECHANIC SETTLEMENT PLUTON (mainly gabbro and diorite)	557 ± 3 (3)	
LOWER COLDBROOK GROUP			
FBft	<b>Fletcher Brook Fm</b> : mainly varicoloured dacitic lapilli tuff and dark grey to black dacitic flows	555.5 ± 1.9 (5)	
WDvs	<b>Walton Dam Formation</b> : varicoloured lapilli tuff and tuffaceous conglomerate with interlayered laminated grey-green siltstone and basalt lenses and layers		551.38 ± 0.24 (NB18-381)
CEfc	<b>Cedar Camp Formation</b> : laminated black to grey to green siliceous siltstone; minor tuffaceous conglomerate		
SBft	<b>Saddleback Brook Formation</b> : dark grey to black (locally red to pink) lapilli and crystal tuff of andesitic to rhyolitic composition, locally flow-banded; minor basaltic lenses		
BENit	<b>Ben Lomond Fm</b> : volcanogenic lapilli tuff or tuffaceous conglomerate, typically with subrounded to subangular andesitic, dacitic, & rhyolitic clasts; locally black dacitic lapilli tuff, grey crystal tuff, and laminated siltstone/chert		
MBiv	<b>McBrien Lake Fm</b> : green-grey mainly dacitic flows and crystal tuff; minor tuffaceous sandstone	557.4 ± 2.5 (5) 554 ± 14/-1 (2)	551.57 ± 0.23 (NB21-451)

**Figure 2.** Summary of main formations and rock types in the lower and upper Coldbrook Group listed in postulated stratigraphic order modified after Barr and White (2004). The Silver Hill Formation is subdivided by area as shown on Figure 1, and the Hosford Brook Formation is inferred to be similar in age to the younger Fundy Trail Parkway rhyolite. Dated plutons are shown above the formations that they intruded based on mapping by Barr and White (1999b, 2004). Previously published U–Pb dates are shown in black in the first column on the right of the figure, with source reference in brackets: (1) Bevier and Barr (1990); (2) Barr *et al.* (1994); (3) Grammatikopoulos *et al.* (1995); (4) Miller *et al.* (2000); (5) Barr *et al.* (2020). Red numbers in the second column are dates and sample numbers from Escribano *et al.* (2023), positioned in approximate stratigraphic position relative to the less precise previously published dates. Some minor formations are omitted from the diagram. Unit colours match Figure 1.

their arc characteristics were interpreted to have been inherited from their source rocks and/or contamination by those rocks during magma evolution (Currie and Eby 1990; Barr and White 1999a). The cause of voluminous but localized magmatism in the Caledonia terrane, and the reason for the evolution from mainly intermediate-felsic to subsequent bimodal magmatism were not considered in those publications.

The present study is based in part on a compilation of previously reported petrographic, whole-rock major and trace element, and Sm–Nd isotopic data from volcanic and plutonic rocks (Guy 1998; Barr and White 1999a; Samson *et al.* 2000). Those historical data are combined with new petrographic, whole-rock geochemical, and Sm–Nd isotopic data to better interpret the characteristics of both the intermediate-felsic and bimodal magmatism and the genesis of these voluminous magmas that led to late Ediacaran super-eruption(s) in the Caledonia terrane.

## REGIONAL GEOLOGY

The Caledonia–Clover Hill Fault separates the Caledonia terrane from adjacent rocks of the Brookville terrane and other Ganderian terranes to the north (Fig. 1 and inset map; Barr and White, 1999a, b, 2004). The southern boundary is inferred to be the extension of the Cobequid–Chedabucto Fault system through the Bay of Fundy, separating Caledonia and the Avalonian terranes in Nova Scotia from the adjacent Meguma terrane to the south (Fig. 1, inset map).

The term Coldbrook Group was originally applied to volcanic rocks throughout southern New Brunswick (e.g., Ruitenberg *et al.* 1979; Dostal and McCutcheon 1990) but based on field observations, geochemistry, and a few widely separated U–Pb zircon dates from volcanic and plutonic samples, Barr and White (1999a, b) divided volcanic rocks in the Caledonian Highlands into an older (ca. 620 Ma) Broad River Group and a younger (ca. 560–550 Ma) Coldbrook Group, and excluded volcanic rocks from Ganderian terranes north of the Caledonia–Clover Hill Fault from those groups. More recent work in the Caledonian Highlands led to the recognition of even older volcanic rocks, now termed Lumsden Group, based on ca. 690 Ma U–Pb zircon dates from cross-cutting plutons (Fig. 1; Barr *et al.* 2019, 2020; 2023). Both the Lumsden and Broad River groups and associated plutons are interpreted to have formed in continental margin arcs (Barr and White 1999a) and are exposed mainly in the eastern and southern highlands (Fig. 1). Although contacts are now mainly tectonic, the Coldbrook Group was likely deposited unconformably on the older units (Park *et al.* 2008; 2017; Barr *et al.* 2020, 2023). The Hammondvale Metamorphic Suite on the northern margin of the Caledonia terrane contains high pressure-low temperature (9 kb, 580°C) pelitic and calcareous rocks that have been interpreted to have been part of an accretionary prism related to the Broad River Group arc (Fig. 1; White *et al.* 2001).

The ca. 550 Ma non-marine clastic sedimentary rocks of the Coldbrook Group are overlain by marine fossiliferous siliciclastic platformal sediments of the Cambrian–Ordovician Saint John Group that are considered typical of Avalonia (Tanoli and Pickerill 1988; Landing 1996; Landing and Westrop 1998; Palacios *et al.* 2011; Alvaro *et al.* 2023).

In the central and western parts of the Caledonian Highlands, both the Coldbrook and Saint John groups are affected by regional folds plunging to the north-northeast (Fig. 1; Barr and White, 1999a, b, 2004). In contrast, in the southeastern part of the highlands, the Coldbrook Group has been thrust over the Lumsden and Broad River groups and related plutons along the Arnold Lake and St. Martins–Stuart Mountain high strain zones, and the younger bimodal units of the Coldbrook Group are mylonitic in the St. Martins–Stuart Mountain high-strain zone (Fig. 1; Park *et al.* 2008, 2017).

## COLDBROOK GROUP STRATIGRAPHY

Barr and White (1999a, b) divided the Coldbrook Group formations into lower and upper based on sparse evidence observed in the field for younging directions and the fact that units in the lower part are intruded by plutons including Bonnell Brook, Baxters Mountain, Upham Mountain, and Mechanic Settlement (Figs. 1, 2). That terminology is retained here, although Escribano *et al.* (2023) showed, using precise geochronology, that ages of the plutons and their host volcanic rocks of the lower Coldbrook Group are similar to the age of rhyolite in part of the Silver Hill Formation in the upper Coldbrook Group and that much of the Coldbrook Group and related plutons formed in less than 760 000 years (Fig. 2). Formations in the lower Coldbrook Group consist predominantly of andesitic to dacitic lithic and/or crystal lapilli tuff and less abundant mafic, intermediate and rhyolitic flows with an estimated thickness of about 10 km (Barr and White 1999a). The intermediate tuffs are characterized by a flow-banded matrix of cryptocrystalline to very fine-grained aggregates of quartz and feldspar. They typically contain sparse (<5%) plagioclase crystals as well as mafic and felsic lithic fragments (Escribano 2021).

Basalt flows are a minor component of the lower Coldbrook Group and mostly consist of clinopyroxene and plagioclase intergrown in intergranular texture with variable amounts of secondary saussurite, epidote, and chlorite (Barr and White 1999a). Interlayered volcanogenic sedimentary formations consist of laminated red to grey siltstone, minor conglomerate, epiclastic sandstone, and laminated grey siltstone, and were described in more detail by Satkoski (2008) and Satkoski *et al.* (2010).

The upper Coldbrook Group formations consist mainly of basalt and rhyolite. Basalt-dominated formations extend from near the city of Saint John to the northeast for about 75 km. The most extensive formation, Hosford Brook, is typically amygdaloidal, with amygdales filled by epidote,

chlorite, calcite, albite, and quartz. The groundmass consists mainly of albitized plagioclase, chlorite, epidote, and opaque phases (Barr and White 1999a). The Burley Lake and Dolan Brook formations include minor interlayered rhyolite and dacite, and in the case of Burley Lake, red conglomerate and sandstone (Fig. 2).

Barr and White (1999a) assigned felsic volcanic rocks in the upper Coldbrook Group to the Silver Hill Formation, which occurs in the Silver Hill area in the northwestern highlands, the Blackall Lake area near Saint John, the Vernon Mountain area, and in a belt that extends from the St. Martins area for 75 km to the northeast parallel to the basaltic Hosford Brook Formation (Fig. 1). The latter belt is informally termed the Fundy Trail Parkway rhyolite (FTPR), and a sample from that unit yielded an age of  $549.18 \pm 0.09$  Ma, 2.5 million years younger than the age of  $551.70 \pm 0.20$  Ma for a rhyolite sample from the Silver Hill area (Fig. 2). Although the rhyolites in the Blackall Lake and Vernon Mountain areas are undated because samples from those areas did not yield zircon (Escribano *et al.* 2023), they are interpreted to be similar in age to rhyolite in the Silver Hill area. Furthermore, the close association between the Hosford Brook Formation and Fundy Trail Parkway rhyolite suggests that it is also younger than the other basaltic units, leading to a revised stratigraphic interpretation (Fig. 2). Rhyolite in the Silver Hill, Vernon Mountain, Blackall Lake areas is typically flow-banded with sparse plagioclase phenocrysts and lithic fragments. The Blackall Lake rhyolite contains highly vesicular pumice fragments, the rhyolite in the Silver Hill area contains lenses with spherulitic textures, and in the Vernon Mountain area, the rhyolite is characterized by perlitic textures and rare lithic clasts with large quartz and feldspar grains and fine aggregates of chlorite and pyrite, rimmed by iron oxide (Escribano 2021). The Fundy Trail Parkway rhyolite is also typically flow-banded with sparse quartz crystals and plagioclase crystals and fragments in a fine-grained groundmass of quartz and feldspar. In places it contains lenses with perlitic fractures and spherulitic texture (Escribano 2021). In the eastern Fundy Trail Parkway area and farther northeast, the rhyolite is moderately to pervasively mylonitized in the St. Martins–Stuart Mountain high-strain zone.

## PLUTONIC ROCKS

The largest pluton is Bonnell Brook (Fig. 1), composed mainly of syeno- and monzogranite that intruded units of the lower Coldbrook Group. The modal mineralogy is plagioclase (35–45%), quartz (35–40%), perthitic orthoclase (30–35%), chloritized biotite (<3% wt.), and rare amphibole. Accessory phases include magnetite, titanite, and less abundant allanite, rutile, apatite, and zircon (Guy 1998; Escribano 2021). Toward the margins of the main body, the medium- to coarse-grained texture in the granite changes to fine-grained allotriomorphic inequigranular to porphyritic, as a result of more rapid cooling (Escribano 2021). The south-

eastern part of the pluton consists of dioritic rocks, locally intruded by granitic dykes (Guy 1998; Barr and White 1999a). The dioritic rocks consist mainly of plagioclase and hornblende, with minor interstitial quartz, K-feldspar and relict clinopyroxene in the cores of some hornblende grains (Barr and White 1999a).

A fine-grained granitic dome with spherulitic to granophyric texture and miarolitic cavities occurs north of the main body of the Bonnell Brook pluton and has been previously interpreted to be a subvolcanic intrusion that is part of the Bonnell Brook pluton and cogenetic with the Silver Hill Formation (Guy 1998; Barr and White 1999a). The samples from this dome consist of plagioclase, quartz, and K-feldspar microphenocrysts in a fine-grained groundmass of quartz and feldspar, and highly chloritized biotite and amphibole (Escribano 2021). Granitic samples from the Upham Mountain pluton farther to the southwest (Fig. 1) also have spherulitic to granophyric textures and miarolitic cavities, and that pluton may also be a subvolcanic intrusion, although it has not yet been mapped in detail.

Other plutons interpreted to be comagmatic with the Coldbrook Group based on field relations, rock types, and previous geochronology include the bimodal Baxters Mountain pluton and the layered mafic-ultramafic Mechanic Settlement pluton (Fig. 1; Barr and White 1999a, b, 2004). The petrology of the Mechanic Settlement pluton and associated Cu-Ni and platinum-group-element mineralization were studied previously (Grammatikopoulos 1992; Grammatikopoulos *et al.* 1995, 2007; Hiebert 2005; Hiebert *et al.* 2008) and hence that pluton is not included in the present study.

## WHOLE-ROCK GEOCHEMISTRY

### Methodology

To investigate the chemical characteristics of the Coldbrook Group and associated plutons, previously acquired whole-rock geochemical data were compiled from Guy (1998) and Barr and White (1999a and unpublished data). The data are from 89 volcanic samples and 68 plutonic samples and were obtained by different methods as documented in the source references. Because the initial work was done decades ago, trace element data, especially rare-earth element (REE) data, were few and of uncertain quality. Hence, 43 of the archived powders were re-analyzed for REE and selected other trace elements by N. Eby at the University of Massachusetts (Lowell) in the early 2000s, using X-ray fluorescence (XRF) and instrumental neutron activation methods as described by Eby and Currie (1996). For the present study, an additional 68 of the archived sample powders were reanalyzed for trace elements by inductively coupled plasma-mass spectrometry (ICP-MS) at Bureau Veritas Lab in Vancouver, BC. For those samples and those analyzed by N. Eby, the existing trace element data in the data compilation

## Silicic samples

were replaced by the new data. In addition, 38 silicic volcanic and plutonic samples were collected as part of the present study to facilitate a more detailed investigation of the relationships among those units. The new samples were also analysed at Bureau Veritas Lab in Vancouver using XRF and ICP-MS methods. For quality control, 5 duplicates, 12 reference materials, and 6 blanks were included in the analytical runs. The analytical data used in the diagrams presented in this paper are tabulated in the M.Sc. thesis of Escribano (2021), available digitally in the Acadia University Library, and in the Supplementary data table that accompanies this paper.

As part of the present study, Sm–Nd isotopic data were obtained from lithic tuff samples NB19-381 and NB19-391 from the lower Coldbrook Group and rhyolite samples NB15-350 and NB19-390 from the upper Coldbrook Group (Table 1). Analyses were done at the Department of Earth Sciences, Memorial University, St. John's, NL, using mass spectrometry (Table 1 footnotes). These new data are compared to previously published Sm–Nd isotopic data from 11 igneous samples from the Broad River Group and related plutons and 12 samples of the Coldbrook Group and related plutons from Whalen *et al.* (1994), Samson (1995), Samson *et al.* (2000), and Barr *et al.* (2000). The software package REFLEX™ from ioGAS™ was used to display the compiled chemical data on diagrams to illustrate their compositional characteristics and tectonic setting.

## Overview

The lower Coldbrook Group samples span a wide range of compositions from mafic to felsic, whereas samples from the upper Coldbrook Group are generally bimodal (Fig. 3a, b, c). The associated plutons are also bimodal, and previous interpretations suggested that the plutons are intrusive equivalents of the basalt and rhyolite in the upper Coldbrook Group (e.g., Barr and White 1999a). The results of more precise U–Pb dating suggest that this interpretation may be appropriate for some rhyolite such as that in the Silver Hills area, but it is not the case for the younger Fundy Trail Parkway (FTP) rhyolite and associated Hosford Brook Formation basalt of the uppermost Coldbrook Group (Fig. 2).

All analyzed samples are subalkalic (Fig. 3b) and the alkalis-FeO<sup>T</sup>-MgO (AFM) diagram shows iron enrichment suggesting that they are tholeiitic (Fig. 3c). Most of the silicic samples from both the lower and upper Coldbrook Group plot in the igneous spectrum for unaltered rocks (Fig. 3d). A few samples, mostly from the lower Coldbrook Group, show potassium enrichment and about one-third of the andesitic and dacitic samples from the lower Coldbrook Group and most of the mafic samples from both the lower and upper Coldbrook Group show a gain in sodium, probably related to albitization (Fig. 3d). Thus, the Coldbrook Group and related plutons are not highly altered and with caution, some mobile elements may be useful to characterize them.

The silicic samples (those with >70% SiO<sub>2</sub>) are from the lower Coldbrook Group, felsic domes and plutons, and the Blackall Lake, Silver Hill, Vernon Mountain, and FTP rhyolites (Fig. 3a). Two trends are visible on the Zr/Y–Y diagram, with the lower Coldbrook Group rhyolite, Blackall Lake and Vernon Mountain rhyolite and felsic dome samples showing higher Zr/Y ratios compared to samples from the felsic plutons and Silver Hill and FTP rhyolites (Fig. 4a). The silicic volcanic and plutonic samples plot mostly in the transitional field, with fewer samples in the tholeiitic field and even fewer in the calc-alkalic field (Fig. 4a).

On the Nb–Y diagram (Fig. 4b), most samples plot in the field for within-plate granite, with a few samples, mostly from plutons, plotting in the field for volcanic-arc granite (Fig. 4b). On the Rb–Hf–Ta ternary diagram of Harris *et al.* (1986), approximately half of the volcanic and felsic dome samples plot in the volcanic-arc field whereas the other samples plot in the field for within-plate granite (Fig. 4c). In contrast, the felsic pluton samples plot mostly in the field for late- and post-collisional granite with fewer samples in the within-plate granite field (Fig. 4c). Most samples from the lower Coldbrook Group, Vernon Mountain and Blackall Lake rhyolites, and felsic domes have relatively high Zr contents and plot in the field for within-plate granite (Fig. 4d). In contrast, most samples from the FTP rhyolite, felsic plutons, and Silver Hill rhyolite have lower Zr and plot in the field for I-, S-, and M-type granite (Fig. 4d). Most of those samples also have Ga/Al ratios within the range typical of I-, S-, and M-type granite, although the FTP rhyolite samples have higher Ga/Al ratios than most other low Zr samples (Fig. 4d), indicating that these rocks are more evolved (Collins *et al.* 1982; Whalen *et al.* 1987).

Chondrite-normalized rare-earth element (REE) plots for the silicic samples of the Coldbrook Group and related plutons show similar patterns, with moderate slopes from light REE to middle REE and flat patterns from middle REE to heavy REE (Fig. 5a–c). The REE concentrations vary between 10- and 300-times chondrite. Average negative Eu anomalies are well developed in every sample, and are probably related to plagioclase fractionation (e.g., Holder *et al.* 2015). The least pronounced Eu anomalies are in the felsic dome samples with average [Eu/Eu\*] of 0.54, and the most pronounced Eu anomalies are in the felsic pluton samples with an average [Eu/Eu\*] of 0.45 (Fig. 5c). Average [La/Sm]<sub>N</sub> ranges from 2.80 to 3.36, with the lowest and highest values in the FTP rhyolite and felsic pluton samples, respectively. Average [Gd/Yb]<sub>N</sub> values range from 1.00 to 1.06. However, the middle REE to heavy REE contents in the felsic pluton samples show a wide range, with contents that vary from 10- to 60-times chondrite, compared to 30- to 60-times chondrite in the felsic dome samples, and 30- to 100-times chondrite in the FTP rhyolite samples. The FTP rhyolite samples show a wide range in light REE and some middle REE, and a narrow range in heavy REE, which

Table 1. New Sm–Nd isotopic data for felsic samples\* from the Coldbrook Group.

SAMPLE	UNIT	Easting	Northing	Lithology	Age (Ma)	$^{143}\text{Nd}$	$^{147}\text{Sm}$	$^{147}\text{Sm}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}$ (t)	$\epsilon\text{Nd}(t)$	TDM (Ma)
NB18-381(2)	Walton Dam Fm	319403	5042156	dacitic tuff	551.38	35.48	8.1	0.13790	0.512478	0.511980	1.1	1152
NB19-390(5)	Silver Hill Fm (FTPR)	326435	5045221	rhyolite	549.18	46.92	10.51	0.13540	0.512475	0.511988	1.2	1124
NB15-350(6)	Silver Hill Fm (FTPR)	318792	5036263	rhyolite	549	36.83	8.14	0.13360	0.512482	0.511999	1.5	1089
NB19-391(7)	Saddleback Brook Fm	318792	5036263	felsic tuff	551	30.73	6.85	0.13480	0.512450	0.511963	0.8	1160

Notes:

\*Numbers in brackets after sample numbers refer to numbered locations shown on Figure 1.

Analyses were done at Memorial University by thermal ionization mass spectrometry using a Finnigan MAT 262V mass spectrometer

(<https://www.mun.ca/creait/the-earth-resources-research-and-analysis-facility/thermal-ionization-mass-spectrometer-facility-tim/>)


Depleted mantle age (TDM) was calculated using a linear evolution for a mantle separated from the CHUR at 4.55 Ga and having a present day epsilon value of +10.

Ratio of  $^{143}\text{Nd}/^{144}\text{Nd}$  is corrected from the deviation from JNdi-1 (mean value of MUN TIMS Lab is 0.512096,  $\sigma = 7$ ,  $n = 16$ ).


Easting and northing coordinates are in NAD83 and grid zone 20.

Ages in blue were obtained by TIMS (thermal ionization mass spectrometry; Escribano *et al.* 2023). The other two ages are estimated based on TIMS ages in similar units.


Abbreviations: Fm, formation; FTPR, Fundy Trail Parkway rhyolite



**Figure 3.** Diagrams illustrating chemical variation in samples from the Coldbrook Group and related plutons. (a)  $\text{SiO}_2$  against  $\text{Zr}/\text{TiO}_2$  with fields after Le Maitre *et al.* (1989) and Hallberg (1984). Diagram is after Stanley (2017). (b)  $\text{Zr}/\text{TiO}_2$  against  $\text{Nb}/\text{Y}$  with fields after Winchester and Floyd (1977). (c)  $\text{Na}_2\text{O}+\text{K}_2\text{O}$ - $\text{FeO}$ - $\text{MgO}$  ternary diagram. Tholeiitic-calc-alkalic dividing line (red) is after Irvine and Baragar (1971). (d)  $\text{Na}_2\text{O}+\text{K}_2\text{O}$  against  $\text{K}_2\text{O} \cdot 100 / (\text{Na}_2\text{O}+\text{K}_2\text{O})$  with fields after Hughes (1972). Abbreviations: UCG, upper Coldbrook Group; LCG, lower Coldbrook Group.



**Figure 4.** Chemical discrimination diagrams for intermediate and felsic samples from the Coldbrook Group and related granitic plutons and domes. (a) Zr/Y against Y with fields after Leshner *et al.* (1986). (b) Nb against Y with fields after Pearce *et al.* (1984). (c) Hf-Rb/30-Ta\*3 ternary plot with fields after Harris *et al.* (1986). (d) Zr against 10000\*(Ga/Al) with fields after Whalen *et al.* (1987). Abbreviations: BL, Blackall Lake area rhyolite; FD, felsic dome; FP, felsic pluton; FTPR, Fundy Trail Parkway rhyolite; LCG, lower Coldbrook Group intermediate and felsic units; SH, Silver Hill area rhyolite; VM, Vernon Mountain area rhyolite. Symbols for rhyolite samples from the Silver Hill area are displayed with thicker outline in (d).



**Figure 5.** Rare earth element (REE) + Y diagrams normalized to C1 chondrite (normalizing values from McDonough and Sun 1995). (a) Fundy Trail Parkway rhyolite, (b) lower Coldbrook Group and Silver Hill, Blackall Lake, and Vernon Mountain area rhyolites, (c) felsic plutons and domes, and (d) upper and lower Coldbrook Group basalt and mafic plutons. Average values per group of samples are shown coloured lines. Average values for  $\text{Eu}/\text{Eu}^*$ ,  $[\text{La}/\text{Sm}]_N$ ,  $[\text{Gd}/\text{Yb}]_N$ , and  $[\text{La}/\text{Yb}]_N$  for each group of samples are shown (normalization values after Taylor and McLennan 1985). Abbreviations: BL, Blackall Lake area rhyolite; FD, felsic domes; FP, felsic plutons; FTPR, Fundy Trail Parkway rhyolite; LCG, lower Coldbrook Group; LCGB, lower Coldbrook Group basalt; MP, mafic plutons; SH, Silver Hill area rhyolite; UCGB, upper Coldbrook Group basalt; VM, Vernon Mountain area rhyolite. In REE diagrams generated by *ioGAS*<sup>TM</sup>, Y is used instead of Tm.


suggests apatite fractionation as apatite has preference for the light REE (e.g., Bruand *et al.* 2020).

Primitive mantle-normalized multi-element plots for the silicic samples show generally flat patterns with trace element contents mainly ranging between 10- and 100-times primitive mantle (Fig. 6a–c). All samples show moderate Nb depletion, and Al contents similar to those of primitive mantle. Scandium contents are variable and generally range between 0.1- and 1-times primitive mantle in the lower Coldbrook Group rhyolite, Silver Hill, Blackall Lake and Vernon Mountain rhyolites, and felsic plutons and domes, whereas in the FTP rhyolite, scandium shows a narrow range between 0.1- and 0.3-times chondrite (Fig. 6), suggesting that neither pyroxene nor hornblende fractionation was involved in the genesis of the younger rhyolite (Wang *et al.* 2021). Vanadium contents show a wide range in the lower Coldbrook Group and Silver Hill, Blackall Lake and Vernon Mountain rhyolites, and are probably related to magnetite

fractionation (Kelley *et al.* 2017), with contents ranging between 0.01- to 0.1-times primitive mantle. Vanadium contents vary between 0.02- to 0.3-times primitive mantle in the felsic plutons and domes.

### Intermediate samples

Samples of andesitic to dacitic composition (56–69%  $\text{SiO}_2$ ) are from the lower Coldbrook Group (Fig 3a). With the exception of two samples with high Zr/Y ratios, they generally have lower Zr/Y ratios compared to the lower Coldbrook Group rhyolite but similar to the Zr/Y ratios of the felsic plutons, FTP rhyolite, and Silver Hill rhyolite (Fig. 4a). They also show lower Nb contents compared to the silicic samples and plot mostly in the volcanic-arc field transitioning to the field for ocean-ridge granite (Fig. 4b). Like the silicic samples, half of the intermediate samples plot in the volcanic-arc field and the other half in the within-plate granite field on the Harris *et al.* (1986) diagram (Fig.



**Figure 6.** Multi-element diagrams with data normalized against primitive mantle (normalizing values from Sun and McDonough 1989). (a) Samples from the Fundy Trail Parkway rhyolite. (b) Samples from the lower Coldbrook Group and Silver Hill, Blackall Lake, and Vernon Mountain area rhyolites. (c) Samples from the felsic plutons and domes. (d) Samples from the upper and lower Coldbrook Group basalt and mafic plutons. Average values per group of samples are represented by coloured lines. Abbreviations: BL, Blackall Lake area rhyolite; FD, felsic domes; FP, felsic plutons; FTPR, Fundy Trail Parkway rhyolite; LC, lower Coldbrook Group; LCB, lower Coldbrook Group basalt; MP, mafic plutons; SH, Silver Hill area rhyolite; UCGB, upper Coldbrook Group basalt; VM, Vernon Mountain area rhyolite.

4c). About two thirds of the samples plot in the field for I-, S-, and M-type granite whereas the remaining samples have higher Zr contents and plot in the field for within-plate granite (Fig. 4d).


Chondrite-normalized REE plots for the andesitic to dacitic samples of the lower Coldbrook Group, like the silicic samples, show moderate slopes from light REE to middle REE and generally flat patterns from middle REE to heavy REE (Fig. 5b). The REE concentrations vary between 20- and 200-times chondrite. Average negative Eu anomalies are 0.73 and  $[La/Sm]_N$  is 2.68. However, the samples show a higher slope in their middle REE to heavy REE contents compared to the silicic samples, as expressed by their higher  $[Gd/Yb]_N$  and  $[La/Yb]_N$  of 1.24 and 4.16, respectively (Fig. 5b).

The andesite and dacite samples of the lower Coldbrook Group show moderate Ti depletion with Ti contents ranging between 3- to 10-times primitive mantle, and a wide range in Sc contents that vary between 0.6- and 2-time primitive mantle (Fig. 6b), suggesting hornblende or pyroxene fractionation. Vanadium contents range between 0.1- to about 2-times primitive mantle in the intermediate samples.

### Basaltic and gabbroic samples

Both the basaltic and gabbroic samples plot in the field for island arc tholeiites on the  $TiO_2$ -MnO- $P_2O_5$  diagram (Fig. 7a). A plot of Th/Yb against Nb/Yb shows that the mafic plutons and lower Coldbrook Group basalt plot above the oceanic array and in the overlapping fields for oceanic and continental arcs, whereas the upper Coldbrook Group basalt samples show a wide range in Th/Yb ratios consistent with a depleted mid-ocean ridge basalt (N-MORB) source contaminated with arc crust (Fig. 7b; Pearce 2014).

On the most recent classification diagram of Shervais (2022), the lower Coldbrook Group basalt and gabbroic pluton samples generally plot between the field for juvenile and mature arcs and continental flood basalts whereas the upper Coldbrook Group basalt samples plot mainly in the field for continental flood basalts (Fig. 7c). Continental flood basalts are characterized by Th/Nb ratios that plot above the oceanic array, and a wide range in Ti/V ratios (20–70 in the Columbia River basalt and 20–43 in the Paraná tholeiites; Shervais 2022).



**Figure 7.** Chemical discrimination diagrams for mafic samples from the Coldbrook Group and related mafic plutons. (a)  $\text{MnO}^*10\text{-TiO}_2\text{-P}_2\text{O}_5$  with fields after Mullen (1983). (b)  $\text{Th}/\text{Yb}$  against  $\text{Nb}/\text{Yb}$  with fields after Pearce (2014). (c)  $\text{Th}/\text{Nb}$  against  $\text{Ti}/\text{V}$  with fields after (Shervais 2022). Abbreviations: DMM, depleted MORB mantle source; EM, enriched MORB mantle source; LCG, lower Coldbrook Group basalt; MORB, mid-ocean ridge basalt; OIB, ocean island basalt; OP, oceanic plateaus; UCG, upper Coldbrook Group basalt. Note: not all mafic samples have trace element data, and hence are not shown on (b) and (c).


Chondrite-normalized REE plots for the mafic samples show smooth patterns with REE concentrations between 10- and 100-times chondrite (Fig. 5d). The REE patterns have gentle slopes from light REE to middle REE, with average  $\text{Eu}/\text{Eu}^*$  ratios of 0.90, 0.90, and 0.87 for the upper and lower Coldbrook Group basalt and gabbroic plutons, respectively (Fig. 5d). Both the upper and lower Coldbrook Group basalt samples have similar light REE contents. The average  $[\text{La}/\text{Sm}]_{\text{N}}$  ratios are 1.91, 1.98, and 2.13 for the upper Coldbrook Group basalt, lower Coldbrook Group basalt, and mafic pluton samples, respectively. The middle REE to heavy REE patterns for the mafic samples are relatively flat with average  $[\text{Gd}/\text{Yb}]_{\text{N}}$  ratios of 1.35, 1.34 and 1.13, respectively (Fig. 5d). However, the lower Coldbrook Group basalt samples have higher average middle and heavy REE contents compared to the upper Coldbrook Group basalt and mafic pluton samples, suggesting a shallower depth of formation for these magmas (e.g., Tegner *et al.* 1998; Melluso *et al.* 2006; Shellnutt *et al.* 2014), and amphibole fractionation expressed by their strong depletion in Y contents (e.g., Dessimoz *et al.* 2012; Fig. 5).

The average  $[\text{La}/\text{Yb}]_{\text{N}}$  ratios for the upper Coldbrook Group basalt, lower Coldbrook Group basalt, and mafic pluton samples are 2.97, 2.45, and 2.59, respectively (Fig. 5d).

Primitive mantle-normalized multi-element plots for the mafic samples show generally flat patterns with trace element contents ranging between 1 and almost 200 times primitive mantle for both basaltic and plutonic samples (Fig. 6d). The mafic samples show moderate Nb and Al depletion, consistent with arc settings and/or contamination by arc lithosphere (e.g., Pearce and Peate 1995), and plagioclase fractionation (McBirney 1984), respectively. They have higher contents of incompatible elements (e.g., Th) than typical MORB (Pearce and Peate 1995), which may be explained by the addition of a lithospheric component. This point is also illustrated in Figures 7b and 7c that show  $\text{Th}/\text{Yb}$  and  $\text{Th}/\text{Nb}$  ratios typical for arc-related rocks.

### Sm–Nd isotopes

The Coldbrook Group and related plutons have  $\epsilon_{\text{Nd}}$  values between -1.5 and +3.5 and model ages between 906 Ma and 1308 Ma (Fig. 8) which are considered typical of Avalonian rocks (e.g., Willner *et al.* 2013; Pollock *et al.* 2015). Also, like older rocks of the Broad River Group and related plutons, which have  $\epsilon_{\text{Nd}}$  values between -0.3 and +4.8 and model ages between 808 Ma and 1422 Ma, most plot in the



**Figure 8 .** Plot of  $\epsilon\text{Nd}$  values against age for samples from the Coldbrook Group and related plutons (coloured symbols) and the ca. 620 Ma Broad River Group and related plutons (black unfilled circles). CHUR: Chondritic uniform reservoir. Data are from this study (Table 1) and Whalen *et al.* (1994), Samson (1995), Samson *et al.* (2000), and Barr *et al.* (2000). Field for West Avalonian isotopic compositions (in grey) is from Murphy and Nance (2002).

Avalonian window of Murphy and Nance (2002; Fig. 8).

The only negative  $\epsilon\text{Nd}$  values are from andesitic and dacitic samples of the lower Coldbrook Group and suggest crustal contamination, whereas all other samples show positive values consistent with mantle-derived melts. The highest  $\epsilon\text{Nd}$  values are displayed by basalt, Vernon Mountain rhyolite, and mafic pluton samples with positive values of 3.5, 3.2, and 2.9, respectively. The FTP rhyolite samples show a narrow range in  $\epsilon\text{Nd}$  values between 1.2 and 1.5, whereas the  $\epsilon\text{Nd}$  values are higher for the felsic pluton samples (0.6 and 1.7) compared to the felsic dome samples (0.4; Fig. 8).

## DISCUSSION

### Magma types

Major and trace element chemistry indicates that the entire late Ediacaran suite, both volcanic and plutonic, is tholeiitic or transitional to tholeiitic. The lower Coldbrook Group andesitic to rhyolitic samples, Silver Hill, Blackall Lake, and Vernon Mountain rhyolites, and felsic dome samples are transitional to tholeiitic, whereas the upper Coldbrook Group, including the FTP rhyolite, and the lower Coldbrook Group basalt and mafic plutons are tholeiitic. However, the multi-element patterns and  $[\text{La}/\text{Yb}]_N$  ratios of 2.6 to 4.4 for the Coldbrook Group and

related plutons are considerably higher than those of typical tholeiitic magmas ( $[\text{La}/\text{Yb}]_N = 1-2$ ), and more within the range for calc-alkalic suites (e.g., Jakes and Gill 1970; Pearce and Peate 1995), although their middle to heavy REE patterns are flatter than those for calc-alkalic magmas, suggesting that they formed in shallower levels in the crust (e.g., Leshner *et al.* 1986), and are consistent with hornblende and zircon fractionation (e.g., Bea 1996).

The Th/Nb and Ti/V ratios for the lower and upper Coldbrook Group basalt and mafic pluton samples are typical for tholeiitic magmas with juvenile arc chemical characteristics (Shervais 2022). Although the upper Coldbrook Group basalts show similar Ti/V ratios as the lower Coldbrook Group basalt and mafic pluton samples, they generally have lower Th/Nb, which suggests that these younger mafic magmas are typical flood basalts with similar trace element contents as those of the Columbia River Group (Shervais 2022).

### Source for magmas

Continental flood basalts (CFB) such as the Columbia River basalt are generally interpreted to have been derived from mixing between upwelling asthenospheric mantle and a large-ion-lithophile-element (LILE)-enriched source (e.g., Wolff *et al.* 2008). The LILE-enriched source is thought to derive from the crust or subcontinental mantle and inherits the chemical characteristics of earlier subduction-related com-

ponents (Hergt *et al.* 1991). Hence, the most crust-contaminated basalt magmas correspond to those that formed the earlier lower Coldbrook Group basalt, whereas the upper Coldbrook Group basalt and mafic plutons are the least contaminated with crustal melts, an interpretation supported by their higher  $\epsilon\text{Nd}$  values.

The mafic samples from the Coldbrook Group and related mafic plutons have low Ti contents which characterizes them as low-Ti CFB generated at relatively shallow depths in the stability field of spinel lherzolite, either in sub-continental lithospheric mantle (SCLM) or in asthenosphere beneath thinned lithosphere (e.g., Tegner *et al.* 1998; Melluso *et al.* 2006; Shellnutt *et al.* 2014). Both models imply the presence of a mantle plume as is commonly linked to the origin of CFB like the Columbia River basalt (e.g., Camp 2013). Hence, we propose that the more crust-contaminated basalt of the lower Coldbrook Group was probably formed from partial melting of sub-continental lithosphere and arc crust, aided by the heat of mantle upwelling, whereas the upper Coldbrook Group basalt (and FTP rhyolite) and mafic plutons was derived from an N-MORB-like source, less contaminated by arc crust, beneath a thinned lithosphere. In the part of Avalonia represented by the Caledonian Highlands, deformation and strike-slip faulting during the Late Ediacaran may have resulted in the opening of a pull-apart caldera (e.g., Acocella *et al.* 2004; Holohan *et al.* 2008), focusing magmatism on thinned lithosphere, however, that process was possibly aided by the presence of a mantle plume, as proposed by Escribano *et al.* (2023) based on zircon chemistry. Other examples of pull-apart calderas occur in Sumatra with the well-known Toba volcano and in the Sierra Madre Occidental in western Mexico (e.g., Ari-bowo 2018; Aguirre-Díaz *et al.* 2021).

#### Lithospheric contamination and calc-alkalic signatures

The generally low Ga/Al ratios of the felsic samples of the Coldbrook Group and related plutons, compared to typical within-plate granite, suggest that they formed from lower temperature liquids (<900°C) derived via partial melting within the subcontinental lithospheric mantle or crust (Pankhurst *et al.* 2011). This interpretation implies that the higher Ga/Al ratios of the FTP rhyolite are a result of less contaminated-higher temperature melts. Zircon chemical data also suggest that the contrasting chemical differences between zircon in samples from the Silver Hill rhyolite, lower Coldbrook Group rhyolite, and felsic domes compared to zircon from the FTP rhyolite reflect higher degrees of crustal assimilation involved in the genesis of the older ca. 551.5 Ma melts, compared to the younger 549.2 Ma rhyolite melt (Escribano *et al.* 2023).

In addition, Nd (this study) and Hf (Pollock *et al.* 2022) isotopic data indicate that samples from the Coldbrook and Broad River groups represent mixing of juvenile magma and isotopically evolved crust at ca. 550 Ma. Pollock *et al.* (2022) further suggested that the vertical trend of  $\epsilon\text{Hf}$  data from ca. 550 Ma samples is best explained by crustal thickening and

recycling of a greater proportion (25%–30%) of an evolved Tonian crustal component. This interpretation supports the suggestion that the lower  $\epsilon\text{Nd}$  values exhibited by samples from the lower Coldbrook Group reflect a larger crustal component compared to samples from the upper Coldbrook Group and related plutons.

#### Chemical differences among felsic products of the 551.5 Ma magmas

Overlapping TIMS U–Pb dates indicate that the lower Coldbrook Group, felsic plutons and domes, Silver Hill area rhyolite, and probably the Vernon Mountain and Blackall Lake rhyolites were contemporaneous and probably derived from a common source. However, they have subtle chemical differences in their high field strength element (HFSE) and REE contents. The high HFSE and other trace element contents of the Blackall Lake and Vernon Mountain rhyolites suggest that those units are closely related to the andesitic to dacitic tuffs of the lower Coldbrook Group and felsic domes. In contrast, the Silver Hill rhyolite has lower HFSE contents than the Vernon Mountain and Blackall Lake rhyolites, lower Coldbrook Group, and felsic dome samples. This difference may be explained by the dripping of solid-liquid mixtures containing hornblende and zircon from the roof of the chamber to the mush below which would gradually lower the HFSE and middle and high REE contents of the overlying melts, represented by the Silver Hill rhyolite.

This mechanism of melt extraction and solid-liquid dripping can also explain the lower HFSE and middle and high REE contents of the felsic plutons compared to the felsic domes, lower Coldbrook Group, and Vernon Mountain and Blackall Lake rhyolite samples, which probably represent the lower mush (e.g., Bachmann and Bergantz 2004) from which the evolved melt was extracted.

#### CONCLUSIONS

The Coldbrook Group and related plutons formed from tholeiitic magmas that originated in an extensional within-plate setting in a region of older continental arc rocks. The basaltic magmas have chemical characteristics of low-Ti continental flood basalts. The more lithosphere-contaminated lower Coldbrook Group magmas were probably derived from older arc lithosphere partially melted by the heat of upwelling mantle. In this model, the melts derived from recycled continental arc lithosphere were of intermediate compositions with arc signatures.

The more evolved Vernon Mountain, Blackall Lake, and Silver Hill area rhyolites represent fractionated melts derived from the lower Coldbrook Group and felsic domes and plutons. The lower HFSE and middle to heavy REE contents of the Silver Hill rhyolite and felsic pluton samples may reflect melt extraction and solid-liquid dripping from the upper part of the magma chamber. Enhanced extension and thinning of the crust, were marked by a change from inter-

mediate to bimodal compositions derived from an N-MORB source, less contaminated with arc lithosphere, that form the youngest basalt and FTP rhyolite of the uppermost Coldbrook Group.

The new data in this study, combined with precise dates reported by Escribano *et al.* (2023), have significantly changed previous ideas of the late Ediacaran magmatic event(s) in the Caledonia terrane and contribute new insights about magmatism during extension in Avalonia.

## ACKNOWLEDGEMENTS

This paper is based in the M.Sc. project of A.P. Escribano. We are grateful to Susan Johnson and Rob Richard of the Geological Surveys Branch (Sussex office) for their help with sample collection when we were unable to travel to New Brunswick due to COVID-19 restrictions in the summer of 2020. Rob Richard also helped in database and map assembly. We thank Chris White for his contributions to mapping of the Caledonia terrane and Adrian Park for his structural insights in the study area. We thank Nelson Eby for giving permission for us to use his trace element analyses of some samples from the Coldbrook Group and Bonnell Brook pluton. We also acknowledge Sherri Strong of Memorial University's CREAT Network for providing Sm–Nd isotopic data. We are grateful to journal reviewers Susan Johnson and Andrea Mills for their helpful comments which led to significant improvements in the manuscript. This work was funded by a research agreement between the Geological Surveys Branch of the New Brunswick Department of Energy and Resource Development and Acadia University, Discovery Grant 2016-04860 from the Natural Sciences and Engineering Research Council of Canada to S.M. Barr, and Acadia University graduate awards to A.P. Escribano.

## URL LINK TO SUPPLEMENTARY DATA

Supplementary Data Table S1: Whole-rock chemical data from the Coldbrook Group and related plutons. <https://journals.lib.unb.ca/index.php/ag/article/view/33598/1882529556>

## REFERENCES

- Acocella, V., Funicello, R., Marotta, E., Orsi, G., and de Vita, S. 2004. The role of extensional structures on experimental calderas and resurgence. *Journal of Volcanology and Geothermal Research*, 129, pp. 199–217. [https://doi.org/10.1016/S0377-0273\(03\)00240-3](https://doi.org/10.1016/S0377-0273(03)00240-3)
- Aguirre-Díaz, G.J., Tristán-González, M., Gutiérrez-Palomares, I., Martí, J., López-Martínez, M., Labarthe-Hernández, G., and Nieto-Obregón, J. 2021. Graben type calderas: The Bolaños case, Sierra Madre Occidental, Mexico. *Journal of Volcanology and Geothermal Research*, 417, 107315. <https://doi.org/10.1016/j.jvolgeores.2021.107315>
- Álvarez, J.J., Johnson, S.C., Barr, S.M., Jensen, S., Palacios, T., Van Rooyen, D., and White, C.E. 2023. Unconformity-bounded rift sequences in Terreneuvian–Miaolingian strata of the Caledonian Highlands, Atlantic Canada. *Geological Society of America Bulletin*, 135, pp. 1225–1242. <https://doi.org/10.1130/B36402.1>
- Aribowo, S. 2018. The geometry of pull-apart basins in the southern part of Sumatran strike-slip fault zone. *IOP Conference Series: Earth and Environmental Science*, 118, 012002. <https://doi.org/10.1088/1755-1315/118/1/012002>
- Bachmann, O. and Bergantz, G.W. 2004. On the origin of crystal-poor rhyolites extracted from batholithic crystal mushes. *Journal of Petrology*, 45, pp. 1565–1582. <https://doi.org/10.1093/petrology/egh019>
- Barr, S.M. and White, C.E. 1999a. Field relations, petrology, and structure of Neoproterozoic rocks in the Caledonia Highlands, southern New Brunswick, Canada. *Geological Survey of Canada Bulletin*, 530, 101 p. <https://doi.org/10.4095/210354>
- Barr, S.M. and White, C.E. 1999b. Geological map of the Caledonia Highlands, southern New Brunswick, Canada. *Geological Survey of Canada Open File* 3615, 4 sheets, scale 1:50 000.
- Barr, S.M. and White, C.E. 2004. Bedrock geology of the Caledonian Highlands of southern New Brunswick. *New Brunswick Department of Natural Resources, Minerals, Policy and Planning Divisions*, Plate 2004-138, scale 1:100 000.
- Barr, S.M., Bevier, M.L., White, C.E., and Doig, R. 1994. Magmatic history of the Avalon terrane of southern New Brunswick, Canada, based on U–Pb (zircon) geochronology. *Journal of Geology* 102, pp. 399–409. <https://doi.org/10.1086/629682>
- Barr, S.M., Hamilton, M.A., White, C.E., and Samson, S.D. 2000. A Late Neoproterozoic age for the Caledonia Mountain Pluton, a high Ti–V layered gabbro in the Caledonia (Avalon) terrane, southern New Brunswick. *Atlantic Geology*, 36, pp. 157–166. <https://doi.org/10.4138/2018>
- Barr, S.M., Johnson, S.C., van Rooyen, D., White, C.E., and Park, A.F. 2019. U–Pb (zircon) dating in the Caledonia Highlands, southern New Brunswick – progress report. *New Brunswick Energy and Resource Department, Information Circular* 2019-1, pp. 10–14.
- Barr, S.M., Johnson, S.C., Dunning, G.R., White, C.E., Park, A.F., Walle, M., and Langille, A. 2020. New Cryogenian, Neoproterozoic, and mid-Paleozoic U–Pb zircon ages from the Caledonia terrane, southern New Brunswick: Better constrained but more complex volcanic stratigraphy. *Atlantic Geology*, 56, pp. 163–187. <https://doi.org/10.4138/atlgeol.2020.007>
- Barr, S.M., Johnson, S.C., van Rooyen, D., Escribano, A.P., Crowley, J.L., and White, C.E. 2023. U–Pb (zircon) dating in the Caledonia Highlands, southern New Brunswick—An update. *In Geological Investigations in New Brunswick*.

- Edited by* E. Smith. New Brunswick Department of Natural Resources and Energy Development; New Brunswick Geological Survey, Geoscience Report 2023-1, pp. 1–29.
- Bea, F. 1996. Residence of REE, Y, Th and U in granites and crustal protoliths; Implications for the chemistry of crustal melts. *Journal of Petrology*, 37, pp. 521–552. <https://doi.org/10.1093/petrology/37.3.521>
- Beranek, L. P., Hutter, A.D., Pearcey, S., James, C., Langor, V., Pike, C., Goudie, D., and Oldham, L. 2023. New evidence for the Baltican cratonic affinity and Tonian to Ediacaran tectonic evolution of West Avalonia in the Avalon Peninsula, Newfoundland, Canada. *Pre-cambrian Research*, 390, 107046. <https://doi.org/10.1016/j.precamres.2023.107046>
- Bevier, M.L. and Barr, S.M. 1990. U–Pb age constraints on the stratigraphy and tectonic history of the Avalon terrane, New Brunswick, Canada. *Journal of Geology*, 98, pp. 53–63. <https://doi.org/10.1086/629374>
- Bruand, E., Fowler, M., Storey, C., Laurent, O., Antoine, C., Guitreau, M., Heilimo, E., and Nebel, O. 2020. Accessory mineral constraints on crustal evolution: elemental fingerprints for magma discrimination. *Geochemical Perspectives Letters*, 13, pp. 7–12. <https://doi.org/10.7185/geochemlet.2006>
- Camp, V.E. 2013. Origin of Columbia River Basalt: Passive rise of shallow mantle, or active upwelling of a deep-mantle plume? *In* The Columbia River Flood Basalt Province. *Edited by* S.P. Reidel, V.E. Camp, M.E. Ross, J.A. Wolff, B.S. Martin, T.L. Tolan, and R.E. Wells. Geological Society of America Special Paper, 497, pp. 181–199. [https://doi.org/10.1130/2013.2497\(07\)](https://doi.org/10.1130/2013.2497(07))
- Collins, W.J., Beams, D., White, J.R., and Chappell, B.W. 1982. Nature and origin of A-type granites with particular reference to southeastern Australia. *Contributions to Mineralogy and Petrology*, 80, pp. 189–200. <https://doi.org/10.1007/BF00374895>
- Currie, K.L. and Eby, G.N. 1990. Geology and geochemistry of the Late Precambrian Coldbrook Group near Saint John, New Brunswick. *Canadian Journal of Earth Sciences*, 27, pp. 1418–1430. <https://doi.org/10.1139/e90-151>
- Dessimoz, M., Müntener, O., and Ulmer, P. 2012. A case for hornblende-dominated fractionation of arc magmas: the Chelan Complex (Washington Cascades). *Contributions to Mineralogy and Petrology*, 163, pp. 567–589. <https://doi.org/10.1007/s00410-011-0685-5>
- Dostal, J. and McCutcheon, S.R. 1990. Geochemistry of Late Proterozoic basaltic rocks from southeastern New Brunswick, Canada. *Precambrian Research*, 47, pp. 83–98. [https://doi.org/10.1016/0301-9268\(90\)90032-L](https://doi.org/10.1016/0301-9268(90)90032-L)
- Eby, G.N. and Currie, K.L. 1996. Geochemistry of the plutons of the Brookville terrane, Saint John, New Brunswick, and implications for development of the Avalon Zone. *Atlantic Geology*, 32, pp. 247–268. <https://doi.org/10.4138/2090>
- Escribano, A.P. 2021. Petrology and age of felsic volcanic rocks of the Coldbrook Group and associated granitic plutons, Caledonia terrane, southern New Brunswick, Canada. Unpublished M.Sc. thesis, Acadia University, Wolfville, Nova Scotia, 263 p.
- Escribano, A.P., Barr, S.M., and Crowley, J.L. 2023. Precise U–Pb zircon dates from silicic super-eruptions during late Ediacaran extension in the Avalonian Caledonia terrane of southern New Brunswick, Canada. *Canadian Journal of Earth Sciences*, 60, pp. 442–462. <https://doi.org/10.1139/cjes-2022-0100>
- Grammatikopoulos, A.L. 1992. Petrogenesis, age, and economic potential of gabbroic plutons in the Avalon terrane in southern New Brunswick and southeastern Cape Breton Island. Unpublished M.Sc. thesis, Acadia University, Wolfville, Nova Scotia, 378 p.
- Grammatikopoulos, A.L., Barr, S.M., Reynolds, P., and Doig, R. 1995. Petrology and age of the Mechanic Settlement Pluton, Avalon terrane, southern New Brunswick. *Canadian Journal of Earth Sciences*, 32, pp. 2147–2158. <https://doi.org/10.1139/e95-167>
- Grammatikopoulos, T.A., Barr, S.M., Hiebert, R.S., Stanley, C.R., and Valeyev, O. 2007. Cu–Ni and PGE minerals in the Mechanic Settlement Pluton, southern New Brunswick, Canada. *The Canadian Mineralogist*, 45, pp. 775–792. <https://doi.org/10.2113/gscanmin.45.4.775>
- Guy, G.H. 1998. Chemical and textural variations in the Bonnell Brook pluton, Caledonian Highlands, southern New Brunswick. Unpublished B.Sc. Honours thesis, Acadia University, Wolfville, Nova Scotia, 131 p.
- Hallberg, J.A. 1984. A geochemical aid to igneous rock type identification in deeply weathered terrain. *Journal of Geochemical Exploration*, 20, pp. 1–8. [https://doi.org/10.1016/0375-6742\(84\)90085-2](https://doi.org/10.1016/0375-6742(84)90085-2)
- Harris, N.B.W., Pearce, J.A., and Tindle, A.G., 1986. Geochemical characteristics of collision-zone magmatism. *In* Collision Tectonics. *Edited by* M.P. Coward and A.C. Rles. Geological Society, London, Special Publications, 19, pp. 67–81. <https://doi.org/10.1144/GSL.SP.1986.019.01.04>
- Hergt, J.M., Peate, D.W., and Hawkesworth, C.J. 1991. The petrogenesis of Mesozoic Gondwana low-Ti flood basalts. *Earth and Planetary Science Letters*, 105, pp. 134–148. [https://doi.org/10.1016/0012-821X\(91\)90126-3](https://doi.org/10.1016/0012-821X(91)90126-3)
- Hiebert, R.S. 2005. Petrogenesis of the Mechanic Settlement Pluton, southern New Brunswick, and controls on associated PGE mineralization. Unpublished M.Sc. thesis, Acadia University, Wolfville, Nova Scotia, 232 p.
- Hiebert, R.S., Barr, S.M., and Stanley, C.R. 2008. Magma evolution and PGE mineralization in the Mechanic Settlement Pluton, southern New Brunswick. *Atlantic Geology*, 44, pp. 78–92. <https://doi.org/10.4138/6507>
- Holder, R.M., Hacker, B.R., Kylander-Clark, A.R.C., and Cottle, J.M. 2015. Monazite trace-element and isotopic signatures of (ultra)high-pressure metamorphism: Examples from the Western Gneiss region, Norway. *Chemical Geology*, 409, pp. 99–111. <https://doi.org/10.1016/j.chemgeo.2015.04.021>

- Holohan, E.P., van Wyk de Vries, B., and Troll, V.R. 2008. Analogue models of caldera collapse in strike-slip tectonic regimes. *Bulletin of Volcanology*, 70, pp. 773–796. <https://doi.org/10.1007/s00445-007-0166-x>
- Hughes, C.J. 1972. Spilites, keratophyres, and the igneous spectrum. *Geological Magazine*, 109, pp. 513–527. <https://doi.org/10.1017/S0016756800042795>
- Irvine, T.N. and Baragar, W.R.A. 1971. A guide to the chemical classification of the common volcanic rocks. *Canadian Journal of Earth Sciences*, 8, pp. 523–548. <https://doi.org/10.1139/e71-055>
- Jakes, O. and Gill, J. 1970. Rare earth elements and the island arc tholeiitic series. *Earth and Planetary Science Letters*, 9, pp. 17–28. [https://doi.org/10.1016/0012-821X\(70\)90018-X](https://doi.org/10.1016/0012-821X(70)90018-X)
- Kelley, K.D., Scott, C.T., Polyak, D.E., and Kimball, B.E. 2017. Vanadium. In *Critical Mineral Resources of the United States—Economic and Environmental Geology and Prospects for Future Supply*. Edited by K.L. Schulz, J.H. DeYoung, Jr., R.R. Seal, and D.C. Bradley. U.S. Geological Survey Professional Paper 1802 pp. U1–U36.
- Landing, E. 1996. Reconstructing the Avalon continent: marginal to inner platform transition in the Cambrian of southern New Brunswick. *Canadian Journal of Earth Sciences*, 33, pp. 1185–1192. <https://doi.org/10.1139/e96-089>
- Landing, E. and Westrop, S.R. 1998. Cambrian faunal sequences and depositional history of Avalonian Newfoundland and New Brunswick: field workshop. In *Avalon 1997 – the Cambrian Standard*. Edited by E. Landing and S.R. Westrop. New York State Museum Bulletin, 492, pp. 5–75.
- Le Maitre, R.W., Bateman, P., Dudek, A.J. and Keller, M.J. 1989. A classification of igneous rocks and glossary of terms, Blackwell, Oxford, 193 p.
- Leshner, C.M., Goodwin, A.M., Campbell, I.H., and Gorton, M.P. 1986. Trace-element geochemistry of ore-associated and barren, felsic metavolcanic rocks in the Superior Province, Canada. *Canadian Journal of Earth Sciences*, 23, pp. 222–237. <https://doi.org/10.1139/e86-025>
- McBirney, A.R. 1984. *Igneous Petrology*. Freeman, Cooper, and Company. p. 270.
- McDonough, W.F. and Sun, S.S. 1995. The composition of the earth. *Chemical Geology*, 120, pp. 223–253. [https://doi.org/10.1016/0009-2541\(94\)00140-4](https://doi.org/10.1016/0009-2541(94)00140-4)
- Melluso, L., Mahoney, J.J., and Dallai, L. 2006. Mantle sources and crustal input as recorded in high-Mg Deccan Traps basalts of Gujarat (India). *Lithos*, 89, pp. 259–274. <https://doi.org/10.1016/j.lithos.2005.12.007>
- Miller, B.V., Barr, S.M., Fyffe, L.R., and White, C.E. 2000. New U–Pb ages from southern New Brunswick: preliminary results. In *Current Research 1999*. Edited by B.M.W. Carroll. New Brunswick Department of Natural Resources and Energy, Minerals and Energy Division, Mineral Resource Report, 2000-4, pp. 39–50.
- Moroz, R. 1994. Petrology and tectonic setting of the Teahan unit and associated base metal mineralization, Caledonian Highlands, southern New Brunswick. Unpublished M.Sc. Thesis, Acadia University, Wolfville, Nova Scotia, 182 p.
- Mullen, E.D. 1983. MnO/TiO<sub>2</sub>/P<sub>2</sub>O<sub>5</sub>: a minor element discriminant for basaltic rocks of oceanic environments and its implications for petrogenesis. *Earth and Planetary Science Letters*, 62, pp. 53–62. [https://doi.org/10.1016/0012-821X\(83\)90070-5](https://doi.org/10.1016/0012-821X(83)90070-5)
- Murphy, B.J. and Nance, D.R. 2002. Sm–Nd isotopic systematics as tectonic tracers: an example from West Avalonia in the Canadian Appalachians. *Earth Science reviews* 59, pp. 77–100. [https://doi.org/10.1016/S0012-8252\(02\)00070-3](https://doi.org/10.1016/S0012-8252(02)00070-3)
- Palacios, T., Jensen, S., Barr, S.M., White, C.E., and Miller, R.F. 2011. New biostratigraphical constraints on the lower Cambrian Ratcliffe Brook Formation, southern New Brunswick, Canada, from organic-walled microfossils. *Stratigraphy*, 8, pp. 45–60.
- Pankhurst, M.J., Schaefer, B.F. and Betts, P.G. 2011. Geodynamics of rapid voluminous felsic magmatism through time. *Lithos*, 123, pp. 92–101. <https://doi.org/10.1016/j.lithos.2010.11.014>
- Park, A.F., Barr, S.M., and White, C.E. 2008. Preliminary investigation of a major high-strain zone in the Caledonian Highlands, southern New Brunswick. *Atlantic Geology*, 44, pp. 127–140. <https://doi.org/10.4138/9717>
- Park, A.F., Treat, R.L., Barr, S.M., White, C.E., Miller, B.V., Reynolds, P.H., and Hamilton, M.A. 2014. Structural setting of the Partridge Island block, southern New Brunswick, Canada: a link to the Cobequid Highlands of northern mainland Nova Scotia. *Canadian Journal of Earth Sciences*, 51, pp. 1–24. <https://doi.org/10.1139/cjes-2013-0120>
- Park, A.F., Barr, S.M., White, C.E., and Johnson, S.C. 2017. The Cambrian to Ordovician Saint John Group east of St. Martins, Fundy Coast Parkway, Saint John County, New Brunswick. New Brunswick Department of Energy and Resource Development, Geoscience Report GR 2017-2, pp. 1–30.
- Pearce, J.A. 2014. Immobile element fingerprinting of ophiolites. *Elements*, 10, pp. 101–108. 316. <https://doi.org/10.2113/gselements.10.2.101>
- Pearce, J.A. and Peate, D.W. 1995. Tectonic implications of the composition of volcanic arc magmas. *Annual Review of Earth and Planetary Sciences*, 23, pp. 251–285. <https://doi.org/10.1146/annurev.ea.23.050195.001343>
- Pearce, J.A., Harris, N.W.B., and Tindle, A.G. 1984. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *Journal of Petrology*, 25, pp. 956–983. <https://doi.org/10.1093/petrology/25.4.956>
- Pollock, J.C., Sylvester, P.J., Barr, S.M., and Murphy, B. 2015. Lu–Hf zircon and Sm–Nd whole rock isotope constraints on the extent of juvenile arc crust in Avalonia: examples from Newfoundland and Nova Scotia, Canada. *Canadian Journal of Earth Sciences*, 52, pp. 1–21. <https://doi.org/10.1139/cjes-2014-0157>
- Pollock, J., Barr, S.M., van Rooyen, D., and White, C.E. 2022. Lu–Hf zircon isotopic constraints on crustal evolution in

- Avalonia and Ganderia in the Canadian Appalachian orogen. *In* New developments in the Appalachian-Caledonian-Variscan orogen. *Edited by* Y. Kuiper, J.B. Murphy, R. D. Nance, R.S. Strachan, and M.D. Thompson. Geological Society of America Special Paper 554, pp. 173–207. [https://doi.org/10.1130/2021.2554\(08\)](https://doi.org/10.1130/2021.2554(08))
- Ruitenbergh, A.A., Giles, P.S., Venugopal, D.V., Buttimer, S.M., McCutcheon, S.R., and Chandra, J. 1979. Geology and mineral deposits, Caledonia area. New Brunswick Department of Natural Resources, Mineral Resources Branch, Memoir 1, 213 p.
- Samson, S.D. 1995. Is the Carolina terrane part of Avalon? *In* Current Perspectives in the Appalachian-Caledonian Orogen. *Edited by* J.P. Hibbard, C.R. van Staal, and P.A. Cawood. Geological Association of Canada, Special Paper 41, pp. 253–264.
- Samson, S.D., Barr, S.M., and White, C.E. 2000. Nd isotopic characteristics of terranes within the Avalon zone, southern New Brunswick. *Canadian Journal of Earth Sciences*, 37, pp. 1039–1052. <https://doi.org/10.1139/e00-015>
- Satkoski, A.M. 2008. Sm–Nd isotopic and whole-rock chemical compositions of late Neoproterozoic and Cambrian sedimentary and metasedimentary rocks of the Caledonian Highlands, southern New Brunswick. Unpublished M.Sc. thesis, Acadia University, Wolfville, Nova Scotia, 150 p.
- Satkoski, A.M., Barr, S.M., and Samson, S.D. 2010. Provenance of Late Neoproterozoic and Cambrian sediments in Avalonia: constraints from detrital zircon ages and Sm–Nd isotopic compositions in southern New Brunswick, Canada. *The Journal of Geology*, 118, pp. 187–200. <https://doi.org/10.1086/649818>
- Shellnutt, J.G., Bhat, G.M., Wang, K.-L., Brookfield, M.E., Jahn, B.-M., and Dostal, J. 2014. Petrogenesis of the flood basalts from the early Permian Panjal Traps, Kashmir, India: geochemical evidence for shallow melting of the mantle. *Lithos*, 204, pp. 159–171. <https://doi.org/10.1016/j.lithos.2014.01.008>
- Shervais, J.W. 2022. The petrogenesis of modern and ophiolitic lavas reconsidered: Ti–V and Nb–Th. *Geoscience Frontiers*, 13, 101319 <https://doi.org/10.1016/j.gsf.2021.101319>
- Stanley, C. 2017. Molar element ratio analysis of lithochemical data: a toolbox for use in mineral exploration and mining. *In* Proceedings of Exploration 17: Sixth Decennial International Conference on Mineral Exploration. *Edited by* V. Tschirhart and M.D. Thomas. Geochemistry Exploration Environment Analysis, 33, pp. 471–494.
- Sun, S.S. and McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. *In* Magmatism in the ocean basins. *Edited by* A.D. Saunders and M.J. Norry, M.J. Geological Society, London, Special Publication, 42, pp. 313–345. <https://doi.org/10.1144/GSL.SP.1989.042.01.19>
- Tanoli, S.K. and Pickerill, R.K. 1988. Lithostratigraphy of the Cambrian–Lower Ordovician Saint John Group, southern New Brunswick. *Canadian Journal of Earth Sciences*, 25, pp. 669–690. <https://doi.org/10.1139/e88-064>
- Taylor, S.R. and McLennan, S.M. 1985. The continental crust: Its composition and evolution. Blackwell, Oxford, 312 p.
- Tegner, C., Leshner, C.E., Larsen, L.M., and Watt, W.S. 1998. Evidence from the rare-earth-element record of mantle melting for cooling of the Tertiary Iceland plume. *Nature*, 395, pp. 591–594. <https://doi.org/10.1038/26956>
- van Staal, C.R., Barr, S.M., McCausland, P.J., Thompson, M.D., and White, C.E. 2021. Tonian–Ediacaran tectonomagmatic evolution of West Avalonia and its Ediacaran–Early Cambrian interactions with Ganderia: an example of terrane transfer due to arc–arc collision? *Geological Society, London, Special Publications*, 503, pp. 143–167. <https://doi.org/10.1144/SP503-2020-23>
- Wang, Z., Yan Hei Li, M., Ray Liu, Z.-R., and Zhou, M.F. 2021. Scandium: Ore deposits, the pivotal role of magmatic enrichment and future exploration. *Ore Geology Reviews*, 128, 103906 <https://doi.org/10.1016/j.oregeorev.2020.103906>
- Whalen J.B., Currie K.L. and Chappell B.W. 1987. A-type granites: geochemical characteristics, discrimination and petrogenesis. *Contributions to Mineralogy and Petrology*, 95, pp. 407–419. <https://doi.org/10.1007/BF00402202>
- Whalen, J.B., Jenner, G.A., Currie, K.L., Barr, S.M., Longstaffe, F.J., and Hegner, E. 1994. Geochemical and isotopic characteristics of granitoids of the Avalon Zone, southern New Brunswick: Possible evidence of repeated delamination events. *Journal of Geology*, 102, pp. 269–282. <https://doi.org/10.1086/629670>
- White, C.E., Barr, S.M., Jamieson, R.A., and Reynolds, P.H. 2001. Neoproterozoic high pressure/low-temperature metamorphic rocks in the Avalon terrane, southern New Brunswick, Canada. *Journal of Metamorphic Petrology*, 19, pp. 519–530. <https://doi.org/10.1046/j.0263-4929.2001.00326.x>
- Willner, A.P., Barr, S.M., Gerdes, A., Massonne, H.-J., and White, C.E. 2013. Origin and evolution of Avalonia: evidence from U–Pb and Lu–Hf isotopes in zircon from the Mira terrane, Canada, and the Stavelot–Venn Massif, Belgium. *Journal of the Geological Society*, 170, pp. 769–784. <https://doi.org/10.1144/jgs2012-152>
- Winchester, J.A. and Floyd, P.A. 1977. Geochemical discrimination of different magma series and their differentiation products using immobile elements. *Chemical Geology*, 20, pp. 325–343. [https://doi.org/10.1016/0009-2541\(77\)90057-2](https://doi.org/10.1016/0009-2541(77)90057-2)
- Wolff, J.A., Ramos, F.C., Hart, G.L., Patterson, J.D. and Brandon, A.D. 2008. Columbia River flood basalts from a centralized crustal magmatic system. *Nature Geoscience*, 1, pp. 177–180. <https://doi.org/10.1038/ngeo124>

*Editorial responsibility: David P. West, Jr.*

## **APPENDIX**

Supplementary data Table S1 is also in open access here:

<https://doi.org/10.25545/XUGV9X>