

Igneous Rock Associations in Canada 3. Large Igneous Provinces (LIPs) in Canada and Adjacent Regions: 3 Ga to Present

R. E. Ernst et K. L. Buchan

Volume 31, numéro 3, septembre 2004

URI : https://id.erudit.org/iderudit/geocan31_3ser01

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)

1911-4850 (numérique)

[Découvrir la revue](#)

Citer cet article


Ernst, R. E. & Buchan, K. L. (2004). Igneous Rock Associations in Canada 3. Large Igneous Provinces (LIPs) in Canada and Adjacent Regions: 3 Ga to Present. *Geoscience Canada*, 31(3), 103–126.

Résumé de l'article

L'histoire de la Terre est ponctuée de nombreuses périodes de mise en place de forts volumes de magma mafiques. De tels magmas qui ne sont pas issus de zones d'expansion « normale » ou de subduction sont appelés Grandes provinces ignées (GPI), et celles-ci sont constituées de basaltes d'épanchements continentaux, de marges de fosse volcaniques, de plateaux océaniques, d'épanchements de basaltes de bassins océaniques, de crêtes sous-marines, et de chaînes de monts sous-marines. Peuvent également y être associées des suites de roches felsiques. Généralement, les GPI du Mésozoïque et du Cénozoïque sont les mieux préservées. Celles du Protérozoïque et du Paléozoïque sont généralement plus fortement érodées et sont constituées de vestiges de basaltes d'épanchement et des réseaux de conduits d'origine (réseaux géants de dykes, provinces de filons-couches et d'intrusifs stratifiées). Dans l'Archéen, les meilleurs candidats sont représentés par les bandes de roches vertes à komatiites. De nombreuses GPI ont été associées à des épisodes de soulèvement régionaux, de dérives ou de fragmentations continentales, ainsi qu'à des crises climatiques. Elles peuvent servir de marqueurs temporels stratigraphiques et sont des cibles de première importance dans l'exploration de gisements de Cu-Ni-ÉGP. Les GPI sont aussi devenues des arguments très considérés dans le débat sur l'existence et la nature des panaches mantelliques.

Le Canada possède de riches archives de GPI, et au moins 80 candidatures ont été isolées sur le territoire canadien et dans les régions adjacentes, leur âge délimitant une fourchette allant de 3 100 Ma à 17 Ma. Nous passons en revue les liens proposés entre la suite des GDI canadiennes d'une part, et celle des panaches mantelliques, des fragmentations continentales, des soulèvements régionaux, et des gisements minéraux, d'autre part. Toutefois, vu le piètre état de caractérisation des unités mafiques au Canada, une campagne de caractérisation géochronologique, paléomagnétique et géochimique serait d'une valeur inestimable pour favoriser l'utilisation des GDI canadiennes pour nous aider à solutionner de grands problèmes géologiques.

SERIES



Igneous Rock Associations in Canada 3. Large Igneous Provinces (LIPs) in Canada and Adjacent Regions: 3 Ga to Present

R.E. Ernst and K.L. Buchan

Geological Survey of Canada

601 Booth Street

Ottawa, Ontario K1A 0E8

ernst@NRCan.gc.ca

SUMMARY

Earth history is punctuated by numerous periods during which large volumes of mafic magma were emplaced. Such magmas not generated by a 'normal' spreading ridge or by subduction are termed Large Igneous Provinces (LIPs), and consist of continental flood basalts, volcanic rifted margins, oceanic plateaus, ocean basin flood basalts, submarine ridges, and seamount chains. Associated felsic rocks may also be present. LIPs of Mesozoic and Cenozoic age are typically the best preserved. Those of Paleozoic and Proterozoic age are usually more deeply eroded, and consist of flood basalt remnants and a deep-level plumbing system (of giant dyke swarms, sill provinces and layered intrusions). In the Archean the most promising LIP candidates are greenstone belts containing komatiites. Many LIPs

have been linked to regional-scale uplift, continental rifting and breakup, and climatic crises. They can be used as precisely dated time markers in the stratigraphic record, and are key targets for Ni-Cu-PGE exploration. LIPs have also become a focus in the debate on the existence and nature of mantle plumes.

Canada has a rich record of LIPs. At least 80 candidates are recognized in Canada and adjacent regions, with ages ranging from 3100 to 17 Ma. We review proposed links between the LIP record of Canada and mantle plumes, continental breakup, regional uplift, and ore deposits. However, given that many mafic units in Canada remain poorly characterized, a concerted geochronology campaign with integrated paleomagnetism and geochemistry would be invaluable in expanding the application of the Canadian LIP record to solving major geological problems.

RÉSUMÉ

L'histoire de la Terre est ponctuée de nombreuses périodes de mise en place de forts volumes de magma mafiques. De tels magmas qui ne sont pas issus de zones d'expansion « normale » ou de subduction sont appelés Grandes provinces ignées (GPI), et celles-ci sont constituées de basaltes d'épanchements continentaux, de marges de fosse volcaniques, de plateaux océaniques, d'épanchements de basaltes de bassins océaniques, de crêtes sous-marines, et de chaînes de monts sous-marines. Peuvent également y être associées des suites de roches felsiques. Généralement, les GPI du Mésozoïque et du Cénozoïque sont les mieux préservées. Celles du Protérozoïque et du Paléozoïque sont généralement plus fortement érodées et sont constituées de vestiges de basaltes d'épanchement et des réseaux de conduits d'origine (réseaux géants de dykes, provinces de filons-couches et d'intrusifs stratifiées).

Dans l'Archéen, les meilleurs candidats sont représentés par les bandes de roches vertes à komatiites. De nombreuses GPI ont été associées à des épisodes de soulèvement régionaux, de dérives ou de fragmentations continentales, ainsi qu'à des crises climatiques. Elles peuvent servir de marqueurs temporels stratigraphiques et sont des cibles de première importance dans l'exploration de gisements de Cu-Ni-ÉGP. Les GPI sont aussi devenues des arguments très considérés dans le débat sur l'existence et la nature des panaches mantelliques.

Le Canada possède de riches archives de GPI, et au moins 80 candidatures ont été isolées sur le territoire canadien et dans les régions adjacentes, leur âge délimitant une fourchette allant de 3 100 Ma à 17 Ma. Nous passons en revue les liens proposés entre la suite des GDI canadiennes d'une part, et celle des panaches mantelliques, des fragmentations continentales, des soulèvements régionaux, et des gisements minéraux, d'autre part. Toutefois, vu le piètre état de caractérisation des unités mafiques au Canada, une campagne de caractérisation géochronologique, paléomagnétique et géochimique serait d'une valeur inestimable pour favoriser l'utilisation des GDI canadiennes pour nous aider à solutionner de grands problèmes géologiques.

INTRODUCTION

Large Igneous Provinces (LIPs) represent voluminous magmatic events that were not generated by a 'normal' spreading ridge or by subduction (Coffin and Eldholm, 1994; 2001; Ernst et al., 2004). They may be emplaced as often as once every 10 Ma (e.g. Coffin and Eldholm, 2001), and time series analysis of the LIP record for the past 3.5 Ga suggests weak cyclicity (Isley and Abbott, 2002; Prokoph et al., 2004).

The most dramatic LIPs are emplaced rapidly (within <10 Ma and often within only a few Ma). These include continental flood basalts, seaward-dipping reflector sequences, oceanic plateaus, and ocean basin flood basalts. Continental flood basalts can be as large as several million cubic km (e.g. the Siberian Traps; Reichow et al., 2002). The largest LIP is the Ontong Java oceanic plateau, which has a volume of 44.4 million cubic km (Coffin and Eldholm, 2001) for combined extrusive (6 million cubic km) and intrusive components (Courtilot and Renne, 2003). The initial large-volume short-duration stage of magmatism of some LIPs has been linked to the arrival of a mantle plume (e.g. White and McKenzie, 1989; Campbell and Griffiths, 1990; Coffin and Eldholm, 1994; 2001; Campbell, 1998, 2001; Ernst and Buchan, 2001; Courtilot et al., 2003). Subsequent rifting/breakup is often associated with a second burst of volcanism (Campbell, 1998) by decompression melting (White and McKenzie, 1989). In addition, LIP magmatism can continue for prolonged periods after the initial outburst (or outbursts), in the form of seamount chains and ridges, which are usually explained as hotspot tracks associated with a plume tail. Other models invoke plate fracturing and 'edge convection' (upper mantle convection between thick and adjacent thin lithosphere), and have been suggested as an alternative to plume models for LIPs and hotspot chains (e.g. Anderson, 2001; Foulger and Natland, 2003).

The volcanic portion of older continental LIPs is largely removed by erosion and deformed during continental collision, whereas older oceanic LIPs are mostly lost during subduction and deformed during ocean closure. Therefore, in the Paleozoic and Proterozoic record, continental LIPs are typically recognized by their exposed plumbing system of giant dyke swarms, sill provinces, large layered intrusions, and remnants of flood basalts (Ernst and Buchan, 2001). The oceanic LIP record may be recognized in some accreted volcanic packages and ophiolite complexes (e.g., Coffin and Eldholm, 2001; Moores, 2002).

The extrapolation of the LIP record into the Archean is more speculative. There are erosional remnants of typical flood basalt provinces, namely the Fortescue sequence of the Pilbara craton of Australia and the Ventersdorp sequence of the Kaapvaal craton of southern Africa (Eriksson et al., 2002). However, most Archean volcanic rocks occur as deformed and fault-fragmented packages termed greenstone belts. Among these, the best candidates for LIPs are thick tholeiite sequences that contain komatiites. The nature of Archean LIPs is discussed in greater detail below.

LIPs are important 1) for testing plume and non-plume models for the generation of LIPs; 2) as precise time markers for stratigraphic correlations; 3) as an aid in reconstructing continents; and 4) as the hosts of major PGE deposits and as a potential tool in diamond exploration. In addition, they can be helpful in studying 5) climatic effects, and 6) regional uplift. We return to these topics after a review of the LIP record of Canada and adjacent regions.

PRELIMINARY LIP HISTORY OF CANADA AND ADJACENT REGIONS

Methodology

Our compilation is based on a recent summary of the global LIP distribution (Ernst and Buchan, 2001) and a newly published compilation of dyke swarms and related magmatic units in Canada and adjacent regions (Buchan and Ernst, 2004). We currently recognize at least 80 LIPs and possible LIP remnants in and adjacent to Canada. The Proterozoic and Phanerozoic mafic magmatic record is reviewed first since its links with LIPs are better defined (Table 1, Fig. 1). The more speculative Archean LIP history follows (Table 2, Fig. 1).

The Proterozoic record relies heavily on diabase dyke swarms and sills (Fig. 2). Dykes injected laterally into the interior of continents have a preservation potential that is much greater than that of associated lavas, and therefore provide a robust record of cratonic LIP events (e.g. Halls, 1982; Fahrig, 1987; Buchan and Halls, 1990).

The compilation includes information on tectonic setting. Our criteria for determining setting rely heavily on dyke swarm geometry and its relationship to cratonic margins (Fig. 3, Table 3). Events are inferred to have a mantle plume origin if a giant radiating dyke swarm is present. Giant linear dyke swarms that extend into a craton (i.e. trend perpendicular to a cratonic margin) are inferred to represent an aulacogen-type swarm ('failed-arm' type in Fahrig, 1987), and can also be used to infer a plume origin with the plume centre situated at the edge of a craton. By contrast, linear swarms that parallel the edge of a craton may simply be rift/breakup related (Ernst and Buchan, 1997) or may possibly represent a back arc rifting setting (e.g. Rivers and Corrigan, 2000), or overriding of a spreading ridge (Gower and Krogh, 2002).

In addition (Table 3), those Archean greenstone belts containing komatiites are inferred to be plume-related on the basis of the elevated temperatures required for generation of komatiites (e.g. Campbell, 1998, 2001; Arndt et al., 1998; Condie, 2001). Finally, small, intraplate events not obviously linked to a cratonic boundary are categorized as 'hotspots'.

Below (and in Tables 1 and 2) we summarize the main events, their age distribution and tectonic setting. It should be noted that referencing has been minimized in the text below because detailed referencing is available through Tables 1 and 2. Also note that we have included number-labels of the form [#14a] in order to facilitate easy cross-correlation with entries in Tables 1 and 2, and with the distribution of main units in Figure 1.

Proterozoic to Present

2.51–2.41 Ga: The earliest Proterozoic LIPs consist of dykes, layered intrusions and volcanic rocks and are mainly associated with the eastern and southern margin of Laurentia. Most notable are the ca. 2.5 Ga Mistassini [#1a] and 2.49–2.45 Ga Matachewan events [#1b] whose radiating diabase dyke swarms locate two plume centres about 800 km apart, and imply rifting of the southeast

Table 1 Large Igneous Provinces (LIPs) and potential LIPs in Canada and adjacent regions since 2.5 Ga. Names of the largest events are underlined. Obsolete names in square brackets. Pre-2.5 Ga record is discussed in Table 2. Anorthosites have not been included. Details and full referencing on most events are available in compilations [1] (=Ernst and Buchan, 2001) and [2] (=Buchan and Ernst, 2004), or in the additional cited references. Abbreviations: se. = southeast, ne. = northeast, c. = central, etc. Units in each entry are ordered in terms of decreasing size. "REF.:" = key reference(s). **SETTING** codes are explained in Table 3.

Event name (Age) (Location) [Size]	Magmatic components (Age)
1 2.51–2.41 Ga	
1a <u>Mistassini</u> (-2.5 Ga) (e. Superior Prov.) [100,000 sq. km]	DYKES: <u>Mistassini</u> (-2.5 Ga) REF: #204 in [1]; #43 in [2] COMMENT: ?Link with <u>Ptarmigan</u> swarm (2.51 Ga [#35 in [2]]) of ne Superior Prov. ?Link with <u>Du Chef</u> swarm (2.41 Ga) [#45 in [2]] along Grenville Front SETTING: Plume (P1); Breakup (B1; se. Superior Craton)
1b <u>Matachewan</u> (2.49–2.45 Ga) (s. & c. Superior Prov.) [360,000 sq. km]	DYKES: <u>Matachewan</u> [<u>Hearst</u>] (2.47 and 2.45 Ga) LAYERED INTRUSIONS: <u>East Bull Lake</u> (2.48 Ga); <u>Agnew</u> (2.49 Ga); <u>River Valley</u> (2.475 Ga); <u>Falconbridge Twp</u> (2.44 Ga) VOLCANIC ROCKS: <u>Dollyberry</u> ; <u>Ellise Mountain</u> ; <u>Copper Cliff</u> rhyolites (-2.45 Ga); <u>Thessalon</u> . REF: #206 in [1]; #42 in [2] COMMENT: ? Link with <u>Ptarmigan</u> swarm (2.51 Ga [#35 in [2]]) of ne. Superior Craton) SETTING: Plume (P1); Breakup (B1; s. Superior Craton)
1c <u>Kaminak</u> (ca. 2.45 Ga (Hearne Prov.) [18,000 sq. km]	DYKES: <u>Kaminak</u> (-2.45 Ga) REF: #203 in [1]; #44 in [2] SETTING: Plume (?P1); ?Breakup (Hearne Craton) COMMENT: Linked with <u>Matachewan</u> event [#1b] by Bleeker (2004)
1d <u>Mirond Lake</u> (2.49 Ga) (<i>Trans Hudson</i>) [two 'windows' into Archean basement about 120 km apart]	DYKES: <u>Mirond Lake</u> (2.49 Ga) REF: #41 in [2] SETTING: ?Breakup (Sask Craton)
2 2.24–2.21 Ga	
2a <u>Kikkertavak</u> (2.235 Ga) (<i>Makkovik Prov.</i>) [6500 sq. km]	DYKES: <u>Kikkertavak</u> (2.235 Ga) REF: #47 in [2] COMMENT: Linked with <u>Lower Aillik Group</u> (2.18 Ga) rift magmatism [Ketchum et al., 2001] SETTING: ?Breakup (s. North Atlantic [Nain] Craton)
2b <u>BN-1</u> (2.21 Ga) (s. <i>Greenland</i>) [54,000 sq. km]	DYKES: <u>BN-1</u> (norite) (2.21 Ga); <u>MD-1</u> undated but possibly similar in age REF: #193 in [1]; #40 in [2] SETTING: ?Breakup (? North Atlantic [Nain] Craton)
2c <u>Ungava</u> (2.22–2.21 Ga) (ne. and e. Superior Prov.) [500,000 sq. km]	DYKES: <u>Ungava</u> radiating swarm = <u>Senneterre</u> (2.22 Ga), <u>Maguire</u> and <u>Klotz</u> (2.21 Ga) SILLS: <u>Nipissing</u> (2.22 Ga) REF: #191 in [1]; #50-52 in [2] SETTING: Plume (P1); Breakup (B1, e. Superior Craton)
2d <u>Malley–MacKay</u> (2.23 and 2.21 Ga) (<i>Slave Prov.</i>) [35,000 sq. km]	DYKES: <u>Malley</u> NE trend (-2.23 Ga); <u>MacKay</u> ENE-E trend (-2.21 Ga) REF: #192 and #197 in [1]; #48 and #57 in [2] SETTING: Breakup (Slave Craton)
3 2.19–2.17 Ga	
3a <u>Dogrib</u> (2.19 Ga) (<i>Slave Prov.</i>) [11,000 sq. km]	DYKES: <u>Dogrib</u> (2.19 Ga) SILLS: <u>Duck Lake</u> (2.18) REF: #185 in [1]; #60 in [2]; Bleeker & Kamo, 2003 SETTING: Breakup (Slave Craton)
3b <u>Tulemalu–MacQuoid</u> (2.19 Ga) (<i>Hearne Prov.</i>) [15,000 sq. km]	DYKES: <u>Tulemalu–MacQuoid</u> (2.19 Ga) REF: #186 in [1]; #61 in [2] SETTING: Breakup (Hearne Craton)
3c <u>Biscotasing</u> (2.17 Ga) (<i>Superior Prov.</i>) [350,000 sq. km]	DYKES: <u>Biscotasing</u> (2.17 Ga) REF: #184 in [1]; #64 in [2]; Halls and Davis, 2004 SETTING: Plume (P3)
3d <u>Cramolet Lake–Payne River</u> (2.17 Ga) (New Quebec Orogen) [30,000 sq. km]	MAGMATISM: 'Cycle 1', including <u>Cramolet Lake</u> sill of basal <u>Seward Subgroup</u> (2.17 Ga) DYKES: <u>Payne River</u> ~2.17 Ga [S. Pehrsson, pers. comm., 2000] REF: #63 in [2] SETTING: Breakup (B2, e. Superior Craton)

Table 1 Continued

4 2.12–2.07 Ga	
4a Marathon (2.12–2.10 Ga) (<i>c. Superior Prov.</i>) [60,000 sq. km]	DYKES: Marathon-N (2.12 Ga); Marathon-R (2.10 Ga) REF: #180 & 181 in [1]; #69 in [2] SETTING: Plume (P2); Breakup (B1, either nc. or sc. Superior Craton)
4b Cauchon Lake (2.09–2.07 Ga) (<i>nw. Superior Prov.</i>)	DYKES: Cauchon Lake (includes Birthday Rapids) (2.09 and 2.07 Ga) REF: #175 in [1]; #72 in [2] SETTING: ?Breakup (B2, nw. Superior Craton)
4c Fort Frances (2.075 Ga) (<i>sw. Superior Prov.</i>) [75,000 sq. km]	DYKES: Fort Frances (2.08 Ga) REF: #174 in [1]; #73 in [2] SETTING: Plume (P2); Breakup (B1, sc. Superior Craton)
4d Lac Esprit (2.07 Ga) (<i>ne. Superior Prov.</i>)	DYKES: Lac Esprit (2.07 Ga) REF: #74-75 in [2], Buchan et al., 2004 SETTING: Plume (P2); ?Breakup (B1, nc. Superior Craton)
4e Griffin (2.11 Ga) (<i>Hearne Prov.</i>) [80,000 sq. km]	SILLS: Griffin [Hurwitz] (2.11 Ga) REF: #179 in [1]; #70 in [2] SETTING: Plume (P3); ?Breakup (Hearne Craton)
4f Napaktok (?2.12 Ga) (<i>Labrador Coast</i>)	DYKES: Napaktok (<2.13 Ga); Tikkigatsiak (2.12 Ga); ? Domes REF: #182 in [1]; #68 in [2] SETTING: ?
5 2.05–2.02 Ga	
5a Kangâmuit (2.05–2.04 Ga) (<i>wc. Greenland; Nain Prov.</i>) [20,000 sq. km]	DYKES: Kangâmuit (of Greenland) (2.04–2.05 Ga); Iglusualiksiuk (2.05 Ga) REF: #170 in [1]; #80 in [2]; [#78 in [2] SETTING: Breakup (?B2, n. North Atlantic [Nain] Craton)
5b Lower Povungnituk (2.04 or 1.96 Ga) (<i>Cape Smith Belt, ne. Superior Prov.</i>)	VOLCANIC ROCKS: Lower Povungnituk SILLS: Korak [2.04 Ga] REF: #169 in [1]; Modeland et al., 2003 SETTING: Breakup (B3, n. Superior Craton)
5c Hearne (2.04 Ga) (<i>Slave Prov.</i>) [15,000 sq. km]	DYKES: Hearne (2.04 Ga) REF: #168 in [1]; #81 in [2] SETTING: Breakup (B2, s. Slave Craton)
5d Lac de Gras (ca 2.025 Ga) (<i>Slave Prov.</i>) [30,000 sq. km]	DYKES: Lac de Gras (2.02–2.03 Ga) LAYERED INTRUSION: Booth River Complex (2.02 Ga) REF: #167 in [1], #82 in [2] SETTING: ?Plume (P3)
6 2.00–1.95 Ga	
6a Minto–Eskimo (2.00 Ga) (<i>e. Hudson Bay, ne. Superior Prov.</i>) [10,000 sq. km for dykes only]	DYKES: Minto (2.00 Ga); Inukjuak VOLCANIC ROCKS: Eskimo; Persillon (Pachi); Nastapoka Group OPHIOLITE: Watts Group (2.00 Ga) REF: #162 in [1]; #88 and #92 in [2]; Scott et al., 1999 COMMENT: geochemically linked with western Povungnituk (2.04 Ga) by Legault et al. (1994) SETTING: Plume (P3); ?Breakup (nc. Superior Craton)
6b Flaherty–Haig (?1.96 Ga) (<i>e. Hudson Bay</i>)	VOLCANIC ROCKS: Flaherty (? -1.96 Ga) SILLS: Haig; Sutton Inlier; Sleeper Island REF: #161 in [1]; Chandler and Schwarz, 1980; Schwarz and Fujiwara, 1981 COMMENT: linked with eastern Povungituk (1.96 Ma) by Legault et al. (1994) SETTING: ?Breakup (B3, nc. Superior Craton)
6c Upper Povungnituk (ca. 1.96 Ga) (<i>Cape Smith Belt, ne. Superior Prov.</i>)	VOLCANIC ROCKS: Upper Povungnituk (1.96 Ga) REF: #169 in [1]; Pehrsson per. comm., 2000; Legault et al., 1994; Modeland et al., 2003 SETTING: Breakup (B3, n. Superior Prov. at 2.04 Ga)
6d Mugford (1.95 Ga) (<i>Nain Prov.</i>) 400 sq. km 1.2 km thick	VOLCANIC ROCKS: Mugford (-1.95 Ga); ? Ramah ; ? Snyder REF: #159 in [1]; Hamilton pers. comm., 2000 SETTING: ? Rifting

Table 1 Continued

7 1.90–1.88 Ga (in Trans Hudson and adjacent Rae-Hearne Craton)	
7a Sandy Bay (1.90 Ga) (<i>Flin Flon belt</i>) (<i>Trans Hudson Belt</i>)	VOLCANIC ROCKS: oceanic plateau (Sandy Bay), oceanic island basalts and ocean floor assemblages REF: #154 in [1]; Syme et al., 1999; Stern et al., 1999 SETTING: ? Plume (P6)
7b Josland (1.88 Ga) (<i>Amisk collage, Trans Hudson</i>)	SILLS: Josland Lake including Mikanagan (1.89–1.88 Ga) REF: Stern et al. 1999; Zwanzig et al., 2001; Turek et al. 2000 SETTING: Back-arc (A2)
7c Bravo Lake (1.90–1.88 Ga) (<i>Piling-Penbryn Group, Rae Prov.</i>)	VOLCANIC ROCKS AND SILLS: Bravo Lake (1.90–1.88 Ga) REF: Jackson and Taylor, 1972; Scott et al., 2002 SETTING: Rift
7d Lake Harbour (-1.87 Ga) (<i>Lake Harbour Group, Rae Prov.</i>)	VOLCANIC ROCKS AND SILLS: Lake Harbour Group (-1.87 Ga) REF: St-Onge et al., 2000 SETTING: Rifting and breakup of the “Meta Incognita microcontinent” from the northern Superior Craton [St-Onge et al., 2000]
8 1.88-1.86 Ga (Circum-Superior events)	
8a New Quebec-Cycle 2 (1.88 Ga) (<i>New Quebec orogen</i>) [30,000 sq. km]	VOLCANIC ROCKS AND SILLS: ‘ Cycle 2 ’ SILLS: Montagnais (1.88–1.87 Ga) REF: #151 in [1] SETTING: Back Arc (A1)
8b Chukotat (1.89–1.87 Ga) (<i>Cape Smith Belt, ne. Superior Prov.</i>)	VOLCANIC ROCKS: Chukotat (lower portion are komatiitic basalts) (1.87 Ga) SILLS: Katniq sills (1.92 or -1.89 Ga) REF: #153 in [1]; N. Wodicka, pers. comm., 2004 SETTING: Plume (P4)
8c Molson–Thompson (1.88–1.86 Ga) (<i>Thompson Belt; nw. Superior Prov.</i>) [>30,000 sq. km]	DYKES: Molson (1.88 Ga); Pickle Crow (1.88 Ga) SILLS: Fox River (1.88 Ga); Thompson Nickel Belt magmatism KOMATIITES: Winnipegosis (1.86 Ga) REF: #150 in [1]; #85 and #98 in [2]; Hulbert et al., 1994; Buchan et al., 2003; Hulbert et al., 2004 SETTING: Plume (?P1, P4); Back-arc (A1)
8d Hemlock (1.88 Ga) (<i>Animikie Basin - Marquette Range, Southern Prov.</i>)	VOLCANIC ROCKS: Hemlock Fm (1.87 Ga); Gunflint Fm. (1.88 Ga); Badwater Greenstone SILLS: Kiernan REF: Ueng et al., 1988; Schneider et al., 2002; Fralick et al., 2002 SETTING: Rifting or Foredeep
9 1.83–1.82 Ga	
9a Sparrow (1.83 Ga) (<i>Rae Prov.</i>) [100,000 sq. km]	DYKES: Sparrow (1.83 Ga) GABBRO BODIES: In Wollaston-Mudjatik Transition zone (1.83–1.82 Ga) REF: #146 in [1] & #101 [2]; Annesley et al., 2003 COMMENT: Is there a link with Christopher Island Fm (Baker Lake basin) (Cousens et al., 2001; Peterson et al., 2002)? SETTING: ? Breakup (Rae Craton)
10 1.75–1.71 Ga	
10a Cleaver (1.75–1.74 Ga) (<i>Bear and Rae Prov.</i>) [three locations spanning an area of 300,000 sq. km]	DYKES: Cleaver (1.74 Ga); Hadley Bay (-1.75 Ga) and MacRae Lake (-1.75 Ga) REF: #105–107 in [2]; Irving et al., 2004. COMMENT: Is this event a precursor to Bonnet River dykes and stocks, and Slab volcanics (1.71 Ga), which have been linked with Wernecke Supergroup (Thorkelson et al., 2001)? Is there a link to Pitz Fm and Nueltin Intrusive Suite of Baker Lake area (Turner et al., 2003; Peterson et al., 2002)? SETTING: ? Plume (P3) Breakup (preceeding breakup of nw. Laurentia)
10b Winagami (?1.89–1.76 Ga) (<i>Western Canada Basin</i>) [120,000 sq. km]	SUBSURFACE SHEETS: Winagami (1.89–1.76 Ga). REF: #149 in [1] COMMENT: Imaged seismically; possible link with Cleaver dykes suggested by Ross and Eaton (1997) SETTING: ? Plume (P3)
11 1.64 Ga	
11a Melville Bugt (ca. 1.64 Ga) (<i>w. Greenland</i>) [220,000 sq. km]	DYKES: Melville Bugt (-1.64 Ga) REF: #132 in [1]; Hamilton et al., 2004 SETTING: ?

Table 1 Continued

12 1.47–1.44 Ga	
12a Moyie (1.47 Ga) (<i>Belt-Purcell basin, Cordillera</i>) [120,000 sq. km dykes only]	SILLS: Moyie (Purcell) (1.47 Ga) VOLCANIC ROCKS: Purcell DYKES: Tobacco Root (Wyoming Block-1) (1.46 Ga) REF: #124 in [1]; #117 in [2] SETTING: Plume (P2); Breakup (w. or sc. Laurentia)
12b Michael–Shabagamo (ca. 1.47 Ga) (<i>ne. Grenville Prov.</i>) [two regions spanning 50,000 sq. km]	SHEETS: Michael Gabbro (-1.47 Ga); Shabagamo (-1.46, 1.45 Ga) REF: Corrigan et al., 2000; Krogh, 1993 <i>in</i> Gower and Krogh, 2002; #122 in [1]; #119-120 in [2] SETTING: Back-arc (A1)
13 1.38 Ga	
13a Hart River–Salmon River Arch (1.38 Ga) (<i>n. Cordillera and s. Belt Basin</i>)	VOLCANIC ROCKS and SILLS: Hart River (1.38 Ga) of n. Cordillera; Salmon River Arch (1.38 Ga) of s. Belt Basin REF: entry in Table 4 of [1]; #121 in [2] SETTING: ? Plume (P2) ? Breakup (nw. Laurentia)
14 1.28–1.27 Ga	
14a Mackenzie (1.27 Ga) (Canadian Shield) [2,700,000 sq. km]	DYKES: Mackenzie (radiating swarm) (1.27 Ga); Bear River (1.27 Ga); '305' VOLCANIC ROCKS: Coppermine; Ekalulia; Nauyat; Hansen; Tweed Lake SILLS: Christie Bay; Tremblay; Goding Bay LAYERED INTRUSIONS: Muskox (1.27 Ga) REF: #107 in [1]; 140, 142, 143 in [2]; Sevigny et al., 1991; French et al., 2002; Schwab et al., 2004 SETTING: Plume (P1); Breakup (B1, n. Laurentia)
14b Harp–Nain–Nutak–middle Gardar 1.28 and 1.27 Ga (North Atlantic Craton) (<i>Nain Prov. and s. Greenland</i>) [80,000 sq. km]	DYKES: Nain-LP (1.28 Ga); Nutak (1.27 Ga); Nain-HP; Harp (1.27 Ga) Gardar-BD0 (-1.28 Ga) Gardar-BD1 REF: #108 in [1]; #132, #133, #137, #138, #139 and #141 in [2] SETTING: ?Breakup (se. Laurentia) COMMENT: The anorthosites, diorites, granites, and troctolites of the 1.35–1.29 Ga Nain Plutonic Suite [e.g. Ryan and James, 2004] which cover an area of 20,000 sq. km in Labrador, are unrelated
15 1.25–1.225 Ga (Grenville Province)	
15a Sudbury [dykes] (1.24 Ga) (<i>Superior and Grenville Prov.</i>) [90,000 sq. km]	DYKES: Sudbury (1.24 Ga) REF: #104 in [1]; #146 in [2] SETTING: Plume (P2); Back-arc (A1)
15b Seal Lake–Mealy (1.25 Ga) (<i>Grenville Prov.</i>) [20,000 sq. km]	VOLCANIC ROCKS: Seal Lake SILLS: Naskaupi (1.25 and 1.22 Ga) DYKES: Mealy (1.25 Ga) North West River. REF: #105 in [1]; #113, #144, & #144a in [2] SETTING: Back-arc (A1)
16 1.18–1.14 Ga	
16a Davy Group–Tshenukutish–Algonquin (1.18–1.16) (<i>Grenville Prov.</i>)	SILLS AND DYKES: Davy Group in Wakeham rift (1.18 Ga); Lillian Suite CORONITIC GABBRO: in Baie du Nord segment of Tshenukutish domain (1.17 Ga) METAGABBROS: Algonquin metagabbros in Central Gneiss Belt (-1.17 Ga) DYKES: Kingston (-1.16 Ga) REF: #97 in [1]; #145, #147 & #149 in [2] COMMENT: Widely separated events in Grenville Prov. of the same age SETTING: Back-arc (A1)
16b Late Gardar (1.16 Ga) (<i>North Atlantic Craton, southern Greenland</i>)	DYKES: Tugtutôq Giant Dykes (1.18 and 1.16 Ga) REF: Buchan et al., 2000; Upton et al., 2003 SETTING: Breakup (?)
16c Abitibi [dykes] (1.14 Ga) (<i>Superior Prov.</i>) [240,000 sq. km]	DYKES: Abitibi (1.14 Ga) REF: #94 in [1]; #155 in [2] SETTING: Plume (P1); Breakup (? precursor to Keweenawan)
17 1.11–1.09 Ga	
17a Keweenawan (Mid-continent rift system) (1.11–1.09 Ga; main pulses: 1.11, 1.10 Ga) (<i>Southern Prov.</i>) [160,000 sq. km]	VOLCANIC ROCKS, SILLS AND DYKES: Keweenawan (1.11–1.09 Ga; main pulses, 1.107 and 1.097 Ga). REF: #90 in [1]; #158-170 in [2] COMMENT: ?Relationship with Southwestern USA Diabase Province [#89 in [1]]; and Moore's Lake sills (1.11 Ga) in Athabasca basin [French et al., 2002]. ?Link with Umkondo event of Africa [#91 in [1]; Hanson et al., 2004] SETTING: Plume (P3) Breakup (attempted breakup of Laurentia)

Table 1 Continued

18 0.78 Ga

18a Gunbarrel (0.78 Ga)
(*Slave Prov.; Mackenzie Mtns. of n. Cordillera; Wyoming Prov.*)
[several areas spanning a distance of about 2500 km]

SHEETS: **Hottah** (sheets) (0.78 Ga)
VOLCANIC ROCKS: **Huckleberry; Irene** (ca. 0.76 Ga)
SILLS: **Faber Lake Gabbro** (0.78); **Wolf Creek** (0.78 Ga); **Mackenzie Mountains DYKES** (0.78 Ga); **MacDonald; Mackenzie Mountains; Tobacco Root-Group B**
REF: #63 in [1]; #178-181 in [2]; Harlan et al., 2003b
COMMENT: ?Link with Windermere rifting [#77 in [1]]; Both South China and Australia have been proposed as the rifted block(s)
SETTING: Plume (P1); Breakup (B1, w. Laurentia)

19 0.72 Ga

19a Franklin-Thule (0.72 Ga)
(*Arctic Islands, mainly Baffin Island; also n. Superior Prov.*)
[1,100,000 sq. km for Canadian portion]

VOLCANIC ROCKS: **Natkusiak**
DYKES: **Franklin** (0.72 Ga); **Thule** (of Greenland)
SILLS: (0.72 Ga) **Minto Inlier; Coronation; Dybbol; Banks Island; Steensby Land** (of Greenland)
REF: #58 in [1] and #185-186 in [2]
COMMENT: Note 0.69 Ga **Gataga** and **Hyland Group** volcanics of northern Cordillera [Ferri et al., 1999; Colpron et al., 2002]. Do these represent subsequent rifting associated with the Franklin-Thule event?
SETTING: Plume (P1); ?Breakup (B1, n. Laurentia)

20 0.615–0.56 Ga (Laurentian margin, Appalachians) **Central Iapetus Event [Puffer 2002]**

20a Long Range (0.62 Ga)
(*e. North America*)
[105,000 sq. km]

DYKES: **Long Range** (0.62 Ga)
VOLCANIC ROCKS: **Lighthouse Cove**
REF: #54 in [1]; #192 in [2]
SETTING: ?Plume ?Breakup (B2, e. Laurentia from Baltica)

20b Grenville-Rideau (0.59 Ga)
(*e. North America*)
[140,000 sq. km]

DYKES: **Grenville** (0.59 Ga)-**Rideau-Adirondack** (fanning swarm)
REF: #53 in [1]; #194–196 in [2]
SETTING: Plume (P1); ?Breakup (B1, e. Laurentia from ?Amazonia)

20c Sept-Îles-Catoctin (0.56 Ga)
(*e. North America*)

LAYERED INTRUSION: **Sept-Îles** (0.56 Ga)
VOLCANIC ROCKS: **Skinner Cove; Catoctin** (in US Appalachians) (0.56 Ga)
REF: #52 in [1]
SETTING: ?Plume (P3); ?Breakup (e. Laurentia from Dashwoods Terrane)

21 0.62–0.55 Ga (Avalon terrane, Appalachians)

21a Harbour Main
(ca. 0.62 Ga)
(*Avalon Zone, Appalachians*)

VOLCANIC ROCKS: part of **Harbour Main Group** (0.63–0.59 Ga)
INTRUSIONS: **Holyrood** granite suite (0.62 Ga); **Connaigre Bay Group** (0.63 Ga); **Simmons Brook suite** (0.62 Ga); **Cap au Miquelon Group** (ca. \geq 6.2 Ga); **East Bay Hills Belt** (ca. 0.62 Ga); **Coxheath Hills Belt** (ca. 0.62 Ga)
REF: Rabu et al., 1996; Bevier et al., 1993
SETTING: ?

21b Marystown
(0.59–0.55 Ga)
(*Avalon Zone, Appalachians*)

VOLCANIC ROCKS: **Marystown Group** (0.59–0.575 Ga); **Long Harbour Group** (0.57–0.55 Ga); **St. Pierre Group** (0.58 Ga); **Belle Riviere Group** (0.57 Ga); part of **Harbour Main Group** (0.63–0.59 Ga); **Coastal Belt** (0.58 Ga)
REF: p. 15, O'Brien et al., 1996; Rabu et al., 1996; Bevier et al., 1993; Fig. 3 in McNamara et al., 2001
SETTING: Arc/ back-arc

22 0.57–0.52 Ga (Cordillera)

22a Hamill-Gog (0.57 Ga)
(*SE Canadian Cordillera*)

VOLCANIC ROCKS: **Hamill-Gog Group** (0.57 Ga)
REF: Colpron et al., 2002
SETTING: ?Breakup

22b Selwyn Basin (0.54–0.45 Ga)
(*Cordillera, n. of and within Selwyn Basin*)

SILLS: **Post-Hyland Group** (0.52 Ga)
VOLCANIC ROCKS: Alkali and potassic mafic volcanism
REF: #218 in [2]; Goodfellow et al., 1995; Abbott, 1997
SETTING: Selwyn Basin rifting

23 0.47–0.42 Ga (Appalachians)

23a Middle Ordovician 'Overstep' Sequence
(0.47–0.45 Ga)
(*Gander Zone-Dunnage Zone*)

VOLCANIC ROCKS: **Middle Ordovician 'Overstep Sequence'** includes bimodal magmatism (in Exploits subzone of Newfoundland, Bathurst area of New Brunswick, Maine, and United Kingdom) (0.47–0.45 Ga)
REF: van Staal et al., 1996
COMMENT: Coeval **Dunn Point** bimodal volc. (0.46 Ga) occur in the Avalon terrane, Nova Scotia (Hamilton and Murphy, 2004)
SETTING: Back-arc

Table 1 Continued

<p>23b Late Ordovician–Silurian magmatism of Atlantic Canada (0.44–0.41 Ga) (c. Newfoundland and Maritimes)</p>	<p>GANDER ZONE–DUNNAGE ZONE (of New Brunswick, Quebec and Maine) (0.43–0.41 Ga) VOLCANIC ROCKS: Tobique and Piscataquis volcanic belts REF: Keppie and Dostal, 1994</p> <p>AVALON ZONE (of Nfld. and New Brunswick) (0.44–0.43 Ga) SILLS: Cape St. Mary's (0.44 Ga) VOLCANIC ROCKS: Bayswater Group (0.44 Ga); Coastal Volcanic Belt (0.43–0.42 Ga) DYKES: Kingston Terrane dyke complex (0.44–0.41 Ga); Passamaquoddy Bay REF: Van Wagoner et al., 2001; Barr et al., 1999; Hodych and Buchan, 1998</p> <p>MEGUMA ZONE (of Nova Scotia) (0.44 Ga) VOLCANIC ROCKS: White Rock Fm (0.44 Ga) INTRUSION: Brenton monzogranite (0.44 Ga) REF: MacDonald et al., 2002; Keppie and Krogh, 2000</p> <p>OVERSTEP SEQUENCE OF NEWFOUNDLAND (0.43–0.42 Ga) VOLCANIC ROCKS: bimodal volc. such as King George IV (0.43 Ga); Springdale (0.42 Ga), and Botwood groups; and mainly felsic volc. such as Sops Arm and La Poile groups INTRUSION: Taylor Brook (0.43 Ga); Main Gut (0.43 Ga) REF: Chandler et al., 1987; Heaman et al., 2002 SETTING: ?</p>
<p>24 0.36–0.32 Ga</p>	
<p>24a Magdalen (Maritimes) Basin (0.36–0.32 Ga) (Maritimes)</p>	<p>INTRUSIONS: magmatic underplate underlying entire Magdalen (Maritimes) Basin, interpreted from gravity; Wentworth and Wyvern gabbros VOLCANIC ROCKS: Cape au Diable; Fountain Lake (mafic to felsic) DYKES: widespread REF: Pe-Piper and Piper, 1998; Murphy et al., 1999 SETTING: Plume</p>
<p>25 0.27–0.20 Ga (Cordillera)</p>	
<p>25a Cache Creek (Late Triassic and mid-Permian) (Cordillera) [~1500 km long, 50–100 km wide]</p>	<p>VOLCANIC ROCKS: Late Triassic volcanics interpreted to represent oceanic plateau (Tardy et al., 2001). Overlies mid-Permian magmatism interpreted to represent underlying Permian Crust Alternative, more complex model suggested by Struik et al. (2001) REF: #38 in [1] SETTING: ?Plume (?P6)</p>
<p>25b Wrangellia (0.23 Ga) (Cordillera) [1,000,000 cu. km]</p>	<p>VOLCANIC ROCKS: Karmutsen; Nikolai INTRUSIONS: Kluane SILLS: Maple Creek (0.23 Ga) REF: #34 in [1]; #303-304 in [2] SETTING: Plume (P6)</p>
<p>25c Ramparts Group (0.21 Ga) (Alaska) [100,000 sq. km]</p>	<p>SILLS AND VOLCANIC ROCKS: Ramparts Group (0.21 Ga) REF: #306 in [2] SETTING: ?</p>
<p>26 0.20 Ga</p>	
<p>26a ENA (Eastern North America) portion of CAMP (Central Atlantic Magmatic Province) (0.20 Ga) (Atlantic Canada) [7,000,000 sq. km for entire event]</p>	<p>VOLCANIC ROCKS: North Mountain Basalt; Grand Manan DYKES: ENA (0.20 Ga) COMMENT: Rest of CAMP event is located along east coast of US and also in Europe, NW Africa and South America REF: #32 in [1]; #307 in [2] SETTING: Plume (P1); Breakup (B1, forming Atlantic Ocean)</p>
<p>27 0.14–0.09 Ga</p>	
<p>27a New England–Québec (NEQ); Monteregean) (0.14–0.11 Ga) [60,000 sq. km]</p>	<p>INTRUSIONS AND PLUTONS: Monteregean Hills intrusions of Quebec and related plutons of New England) (0.14–0.11 Ga) REF: #323 in [2] SETTING: Hotspot</p>
<p>27b Trap (sw. Greenland) [20,000 sq. km]</p>	<p>DYKES: Trap (0.14 Ga) REF: #30 in [1]; #322 in [2] SETTING: ? Breakup (B2, Linked to subsequent breakup of Greenland from Laurentia to form Labrador Sea)</p>

Table 1 Continued

<p>27c Sverdrup Basin Magmatic Province (part of High Arctic Large Igneous Province; HALIP) (0.13–0.09 Ga) (Arctic Islands) [550,000 sq. km]</p>	<p>VOLCANIC ROCKS AND SILLS: including Strand Fiord Fm (0.13–0.09 Ga; major pulse at 0.095 Ga); Alpha Ridge DYKES: Queen Elizabeth Islands (includes Hazen Strait); Lightfoot River INTRUSIONS: Wootton (0.092 Ga) REF: #12 in [1]; #325 in [2]; Tarduno et al., 1998; Trettin and Parrish, 1987 COMMENT: Part of Large Igneous Province that includes Svalbard and Franz Josef Land (Maher, 2001) SETTING: Plume (P1); Breakup</p>
28 0.07–0.05 Ga	
<p>28a Carmacks (n. Cordillera) [63,000 sq. km]</p>	<p>VOLCANIC ROCKS: Carmacks (0.07 Ga) REF: #9 in [1] SETTING: Plume (? P6)</p>
<p>28b Crescent (0.06–0.05 Ga) (western margin of North America)</p>	<p>VOLCANIC ROCKS: Crescent Terrane (“Coast Range Basalt Province”) (0.06–0.05 Ga) REF: #339 in [2]; Murphy et al., 2003 SETTING: Plume (? P6)</p>
<p>28c North Atlantic Igneous Province (NAIP) (0.06 Ga) (Baffin Island, w. Greenland) [1,300,000 sq. km for entire NAIP event]</p>	<p>VOLCANIC ROCKS: Cape Dyer; West Greenland DYKES: Cape Searle; West Greenland REF: #334 and 335 in [2]; #5 in [1] COMMENT: Part of North Atlantic Igneous Province of Greenland, United Kingdom (0.062–0.052 Ga), and linked to present-day Iceland hotspot SETTING: Plume Breakup (Europe from Greenland)</p>
29 0.025–0.015 Ga	
<p>29a Behm Canal (Tertiary ‘Lamprophyre’ Province) (ca. 0.023–0.005 Ga) (c. British Columbia) [25,000 sq. km]</p>	<p>DYKES: Behm Canal (0.023–0.005 Ga) REF: #348 in [2] COMMENT: alkalic lamprophyres SETTING: Hotspot (H1)</p>
<p>29b Columbia River Basalt Group (mainly 0.017–0.015 Ga) (nw. US) [165,000 sq. km]</p>	<p>VOLCANIC ROCKS, DYKES AND SILLS: Columbia River Basalt Group (CRBG) REF: #1 in [1] SETTING: Plume (P1, P7)</p>
<p>29c Chilcotin (Neogene-Paleogene; mainly Miocene) (c. and sc. British Columbia) [25,000 sq. km]</p>	<p>VOLCANIC ROCKS: Chilcotin Group basalts; Anahim volcanic belt; Cheslatta Lake suite; Masset Fm REF: Anderson et al., 2001 COMMENT: Nearby to and potentially linked to Columbia River Basalt Group SETTING: Hotspot (H1) Back-arc (partial melting)</p>

margin of Laurentia at this time. The conjugate margin may have been Baltica, which has a remarkably similar age range of magmatism (e.g. Heaman, 1997; Buchan et al., 2000), or the Hearne craton (Bleeker, 2004). Additional units in eastern and southwestern Laurentia, the 2.505 Ga Ptarmigan dykes [#1a, unit not displayed in Fig. 1] and the 2.408 Ga Du Chef dykes [#1a, unit not displayed in Fig. 1] have uncertain relationship with the events in the southeastern and southern parts of the Superior Province.

Elsewhere in Canada, there are additional events falling in this age range. These include the Kaminak dykes [#1c] in the Hearne Province, and the Mirond Lake dykes [#1d] in the Sask craton of the Trans Hudson orogen.

2.24–2.21 Ga: The next burst of activity in Laurentia was widespread in

the North Atlantic (Nain), Superior, and Slave cratons [#2a–d]. The 2.235 Ga Kikkertavak swarm [#2a] has been linked with breakup along the southern margin of the North Atlantic Craton (lower Aillik Group) at 2.178 Ga (Ketchum et al., 2001) although the long interval between emplacement of the swarm and breakup 57 Ma later is problematic. The 2.21 Ga BN1 (and possibly correlative MD1) dykes [#2b] of western Greenland are also likely linked with North Atlantic Craton breakup.

Superior Province elements of this age include the 2.217–2.210 Ga Ungava giant radiating swarm [#2c] that spans the entire eastern half of the Superior Province (Buchan et al. 1998). The convergence point marks a plume centre and possible breakup of a continental fragment from the eastern

margin of the Superior Province. Buchan et al. (1998) proposed that the Nipissing sills [#2c] of the Southern Province were fed laterally via the Ungava radiating swarm from the plume centre region >1000 km to the north-east.

The Slave Province is another node of activity in this age interval. The 2.23 Ga Malley and 2.21 Ga MacKay dyke swarms [#2d] are both roughly linear, and crosscut at a shallow angle. If they represent rift-parallel swarms (Fig. 3c) then they could be linked to breakup along the southeastern margin of the Slave Province (LeCheminant et al., 1996). However, if they represent an aulacogen (failed-arm) type swarm (Fig. 3b), then the plume centre and locus of breakup would have to lie at one end of the swarms; i.e., either on the east or southwest margins of the Slave Province.




Figure 1 Distribution of mafic units discussed in the text. Numbers correspond to entries in Tables 1 and 2. Colours have no age significance; they are simply used to allow different magmatic events to be distinguished.

Table 2 Archean greenstone belts in Canada interpreted to have a plume origin (i.e. tholeiitic sequences containing komatiites and/ or those interpreted as oceanic plateaus). Details and full referencing on most events is available in compilations [1] (=Ernst and Buchan 2001) and [2] (=Buchan and Ernst 2004), and Tomlinson and Condie (2001), or in the additional cited references. “REF.” = key reference(s). **SETTING** codes are explained in Table 3.

Event name (Age) (Location)	Magmatic components (Age)
A1 3.11–2.98 Ga	
A1a Hunt River (3.11 Ga) (Hopedale block, North Atlantic Craton)	GREENSTONE BELTS: Hunt River (3.11 Ga) REF.: James et al., 2002 COMMENT: Includes komatiites SETTING: ?Plume (P4)
A1b Florence Lake (2.99–2.98 Ga) (Hopedale block, North Atlantic Craton)	GREENSTONE BELTS: Florence Lake (2.99–2.98 Ga) REF.: James et al., 2002 COMMENT: Includes komatiites SETTING: ?Plume (P4)
A1c North Caribou (2.99 Ga) (n. Caribou Terrane, nw. Superior Prov.)	GREENSTONE BELTS: Balmer Assemblage (Red Lake greenstone belt) (2.99 Ga) REF.: #256 in [2]; Tomlinson et al., 1998; Hollings et al., 1999 “plume related rifting” COMMENT: Includes komatiites SETTING: ?Plume (P4)
A2 2.93–2.92 Ga	
A2a Western Superior (2.93 Ga) (Wabigoon subprov., w. Superior Prov.)	GREENSTONE BELTS: Steep Rock; Finlayson; North Rim Unit of McGruer Assemblage (2.93 Ga) Lumby Lake 2963–2898 Ma, ? Heaven Lake [U 2954 Ma] REF.: #252 in [1] COMMENT: Includes komatiites. Rifting or impingement of plume in subduction setting (Hollings et al. 1999) SETTING: ?Plume (P4) Back-arc (A2)
A3 2.86 Ga	
A3a Pickle Crow (2.86 Ga) (Uchi subprov., nw. Superior Prov.)	GREENSTONE BELTS: Pickle Crow Assemblage (-2.86 Ga) REF.: #244 in [1] COMMENT: Includes komatiites SETTING: ?Plume (P4)
A4 2.79–2.78 Ga	
A4a Vizien (2.79 Ga) (Minto Block, ne. Superior Prov.)	GREENSTONE BELTS: Vizien (2.79 Ga) REF.: #236 in [1] COMMENT: Includes komatiites SETTING: ?Plume (P4, P6)
A4b Faribault–Thury (ca. 2.88–2.71 Ga) (Minto Block, ne. Superior Prov.)	GREENSTONE BELTS: greenstone belt remnants in Faribault–Thury Complex (-2.88–2.71 Ga) REF.: Maurice et al., 2003 SETTING: ?Plume (P4)
A4c Fourbay (2.78 Ga) (Wabigoon subprov., w. Superior Prov.)	GREENSTONE BELTS: Fourbay Sequence in Savant Lake and Sturgeon Lake (2.78 Ga) REF.: Sanborn-Barrie and Skulski, 1999 COMMENT: Possible oceanic plateau setting SETTING: ?Plume (P6)
A5 2.75–2.70 Ga	
A5a Prince Albert–Woodburn (2.73 Ga) (Rae Prov.) >1000 km. long belt]	GREENSTONE BELTS: (-2.73 Ga): Prince Albert Group; Woodburn Group; Mary River Group, ?Murmac Bay Group REF.: #220 in [1]; Schau, 1997; Aspler et al., 1999; Zaleski et al., 2001; Hartlaub et al., 2002; Skulski et al., 003a,b. COMMENT: Includes komatiites SETTING: ?Plume (P4) ?Breakup Rae Craton
A5b Kam (2.73–2.70 Ga) (Slave Prov.)	VOLCANIC ROCKS and DYKES: Kam Group and early dykes (2.73–2.70 Ga). REF.: #227 in [1] COMMENT: Widespread tholeiitic cover sequence SETTING: ?Plume (?P3)
A5c Abitibi (2.75–2.70 Ga) (Abitibi belt, Superior Prov.)	GREENSTONE BELTS: includes komatiites: Pacaud (2750–2735 Ma); Stoughton-Roquemaure (2725–2720 Ma); Kidd-Munro (2718–2710 Ma); Tisdale (2710–2703 Ma) REF.: #224 in [1]; Ayer et al., 2002; Sproule et al., 2002 COMMENT: Includes komatiites SETTING: ?Plume (P4); Back-arc (A2)
A5d Wawa (2.75–2.74 Ga) (Wawa belt, Superior Prov.)	GREENSTONE BELTS: Schreiber-Hemlo-White River-Dayohessarah (2.75–2.74 Ga) REF.: #230 in [1] COMMENT: Includes komatiites SETTING: ?Plume (P4)





Figure 2a Aerial photograph (centred approximately at 67°18'N, 112°26'W) of 1.27 Ga Mackenzie dykes cutting the Recluse Group and other units of the Tree River fold belt, Wopmay Orogen. The dykes are up to 50 m wide and are exposed as prominent ridges because of differential erosion.



Printed by GEO. F. DESSABATE. From a Sketch by DR. BELL.
VIEW OF THE OUTER AND INNER "BARNs," LAKE NIPIGON, FROM THE MOUTH OF THE WABINOSH RIVER.
 COLUMNAR TRAP, RESTING ON DOLOMITES, QUARTZITES AND SHALES.

Figure 2b The 1.109 Ga Logan diabase sills of the Keweenawan magmatic event, shown here from an 1869 sketch by Robert Bell of the Geological Survey of Canada (Bell 1879), occur widely in the vicinity of Lake Superior and Lake Nipigon. Bell (1870) noted that "... the Inner Barn with its sides of columnar trap, rises like a great castle in the middle of the bay, to a height upwards of 600 feet, and appears to be the highest point around Lake Nipigon."




Figure 3 LIP setting determined from dyke swarm geometry and relationship to cratonic edge. These and other settings are described in Table 1. Star locates mantle plume center. **a.** Dyke swarm radiating from coeval cratonic margin (focus identifies plume centre) Type P1 and B1 of Table 3. **b.** Linear swarm perpendicular to nearby coeval cratonic margin (failed arm type of Fahrigr, 1987). If a swarm extends across the craton to another margin, it is ambiguous as to which margin the dykes are related. Type P2 of Table 3. **c.** Linear swarm parallel to coeval breakup margin (passive margin type of Fahrigr, 1987). Type B2 of Table 3. **d.** Linear swarm parallel to subducting margin. Type A1 of Table 3.

Table 3 Selected criteria for interpreting origin and setting of LIPs and smaller intraplate mafic events.

Label	Criteria
MANTLE PLUME LINK	
P1	Radiating dyke swarm
P2	Linear dyke swarm (or belt of mafic magmatism) perpendicular to cratonic margin (failed-arm type of Fahrigr, 1987; could also be called aulacogen-type).
P3	Large event, distal from cratonic boundary
P4	Presence of high-Mg rocks, e.g. komatiites in Archean greenstone belts
P5	Geochemistry: OIB or FOZO signature
P6	Accreted oceanic plateaus
P7	Link to present-day hotspot
BREAKUP LINK	
B1	Radiating swarm or failed-arm type swarm (Fahrigr, 1987) with convergence point near cratonic margin of similar age
B2	Linear dyke swarm, parallel to and nearby cratonic margin of similar age. This is termed passive margin type swarm by Fahrigr (1987)
B3	LIP situated nearby cratonic margin of similar age
ARC LINK	
A1	Location in a back-arc setting relative to concurrent orogenic activity. However, this does not necessarily require back-arc extension processes. (See Note 1, below)
A2	Coexisting with arc magmatism
HOTSPOT (OF UNSPECIFIED ORIGIN)	
H1	Small (sub-LIP scale) region of anomalous intraplate mafic volcanism

Note 1: The Keweenawan rift activity is arguably in a back-arc setting with respect to the Grenville orogen, yet it is generally viewed that the Keweenawan magmatism is plume-related (see discussion in text).

2.19–2.17 Ga: Widespread activity occurred during this age interval. The oldest activity is represented by the 2.19 Ga Dogrib dyke swarm [#3a] in the Slave Province and the similar age Tulemalu-MacQuoid swarm [#3b] in the Hearne Province. While a connection would seem likely based on the age match, preliminary paleomagnetic data indicate that the Slave and Rae cratons were not in their present relative positions at this time (LeCheminant et al., 1997). Globally, a stage of Birimian magmatic activity in Western Africa has this age (event #187 in Ernst and Buchan, 2001). Magmatic activity in the Slave Craton continues to 2.18 Ga, the age of the Duck Lake sill [#3a, unit not displayed in Fig. 1].

The 2.17 Ga Biscotasing dyke swarm [#3c] is one of the most widespread in Canada, extending throughout much of the southern and central Superior Craton. Similar age activity [#3d] is present as the Payne River dykes in northern Ungava (Cape Smith Belt) and Cramolet Lake sills (part of "Cycle 1" magmatism) in the Labrador Trough. It seems unlikely that all these elements are part of a single event, but the association of Payne River dykes and "Cycle 1" magmatism in the Labrador Trough suggests a link to breakup along the eastern Superior Province. Globally, an identical age has also been obtained from a quartz diorite dyke in the Wyoming Province, although paleomagnetic data are inconclusive regarding the relative locations of the Wyoming and Superior provinces at this time (Harlan et al., 2003a).

2.12–2.07 Ga: In the Superior Province there are several distinct stages of activity during this time interval. The 2.12–2.10 Ga Marathon dyke swarm [#4a] is a broadly linear swarm that cuts northward across the Superior Province from Lake Superior. This is an aulacogen-type swarm, and therefore could be associated with a breakup margin at the end of the swarm. Since the swarm may reach the cratonic margin on its north end as well as its south end (c.f. Fig. 3b), then the applicable breakup margin may also be at either end, either to the north in Hudson Bay or to the south in Lake Superior.

Slightly younger is the 2.09–2.07 Ga Cauchon swarm [#4b], which has been linked with breakup along the northwest margin of the Superior Province (e.g. Halls and Heaman, 2000). Two other events are also of this age: the 2.077 Ga Fort Frances (Kenora-Kabetogama) dykes [#4c] that are linked with breakup along the southern margin of the Superior Province, and the 2.069 Ga Lac Esprit swarm [#4d] located east of James Bay. There are no obvious connections between these widely separated but coeval 2.07 Ga events. However, future identification of additional dykes of this age in intervening areas could suggest a link.

In the Hearne Province, the widespread 2.111 Ga Griffin gabbros (formerly Hurwitz gabbros) [#4e] are presumably linked to breakup along the Trans-Hudson margin (Aspler et al., 2002).

Finally, the Napaktok swarm [#4f] trends perpendicular to and extends along the Labrador coast for at least 200 km. These dykes have an uncertain age between 2.50 and 2.10 Ga, and uncertain tectonic setting. Although more than one age of dykes may be present, at least part of this activity is dated by the 2.121 Ga age for the Tikkigatsiak dyke.

2.05–2.02 Ga: In the Nain Province, the 2.045 Ga Iglusuatliksuak dyke [#5a, unit not displayed in Fig. 1] is coeval with the Kangâmut swarm [#5a] of adjacent southern Greenland. The combined 2.04–2.05 Ga event would be areally significant, and may represent a breakup event associated with the northern margin of the North Atlantic Craton. Korak sills associated with the lower Povungnituk sequence [#5b] in the Cape Smith Belt of the northeastern Superior Province are of the same age as the Iglusuatliksuak dyke. However, given that the North Atlantic Craton had not yet docked at this time, two independent events are probable. The younger Upper Povungnituk sequence [#6c] is discussed below in the next section.

The third locus of activity of this age is in the Slave Craton. The 2.038 Ga Hearne dykes [#5c] parallel the southern Slave margin, possibly representing breakup along that margin (c.f.

with evidence discussed above for an earlier 2.23–2.21 Ga breakup along that margin associated with the Malley/MacKay swarms [#2d]). In contrast, the younger 2.030–2.023 Ga Lac de Gras swarm [#5d] is not easily linked to any margin. It is centred in the Slave province and slightly converges to the north toward the similar-aged Booth River Complex in the Kilohigok sedimentary basin. LeCheminant et al. (1996) suggest that the Lac de Gras swarm is coeval with rifting on the western margin of the Slave Province.

2.00–1.95 Ga: Magmatism of this age range is distributed along the circum-Superior margin (e.g. Baragar and Scoates 1987).

In the Belcher Islands there are two main volcanic suites, the Eskimo [#6a] and overlying Flaherty [#6b] volcanics. Based on paleomagnetic correlations, the Eskimo volcanics may be linked with Richmond Gulf, Persillon, Pachi and Nastapoka Group volcanics (Chandler and Schwarz, 1980; Schwarz and Fujiwara, 1981), and with 1.998 Ga Minto dykes (Buchan et al., 1998). The Minto dykes and correlated suites are also coeval and perhaps cogenetic with the precisely dated Watts Group ophiolite. The Eskimo volcanics have been linked geochemically with the western Povungnituk [#6c] (Legault et al., 1994), but this link is now doubtful based on more extensive recent geochemical study (Modeland et al., 2003).

The overlying Flaherty Group [#6b] is essentially undated except for a very uncertain Pb-Pb age of 1960±80 Ma. However, paleomagnetic data suggest a link with the Haig and Sutton Inlier sills (Schwarz and Fujiwara, 1981). Geochemical correlations indicate that the Flaherty Group may be linked with the upper (eastern) Povungnituk [#6c] (Legault et al., 1994).

The Cape Smith Belt includes the tectonically juxtaposed 1.998 Ga Watts Group, 2.04–1.96 Ga Povungnituk suites [#5b & #6c], and ca. 1.87 Ga Chukotat Group volcanics [#8b]. (Note that the older 2.04 Ga portion of the Povungnituk sequence [#5b] was discussed in the previous section.)

Additional magmatism in this age interval is present in the Nain Province. A 1.95 Ga plateau basalt sequence in the Mugford Mountains of Labrador [#6d] represents a rifting event (Wardle et al., 2002) that may be more widespread in the North Atlantic Craton based on possible correlations with the Ramah and Snyder groups.

1.90–1.88 Ga (Trans Hudson Orogen and Rae-Hearne Craton):

Several non-arc packages are associated with the Trans Hudson during this age interval. These include the Sandy Bay assemblage of the Flin Flon Belt [#7a], which contains accreted oceanic crust of various affinities, including an oceanic plateau. In addition there are the Josland and related sills [#7b] in the Lynn Lake Belt of the Amisk collage, and the mafic magmatism in the Piling-Penhryn [#7c], and Lake Harbour [#7d] groups on Baffin Island.

1.88–1.86 Ga (Circum-Superior Province): 1.88–1.87 Ga mafic magmatism surrounds the Superior Province on all sides except the southeast where, if originally present, it has been obscured by the Grenville allochthon. This magmatism is present in the Labrador Trough (Montagnais sills belonging to New Quebec Orogen Cycle 2 [#8a]), Cape Smith Belt (Chukotat Group [#8b]), northwest Superior province (Fox River sill, Molson dykes, Thompson Nickel Belt, and Winnipegosis komatiites [#8c]), and the Animikie Basin and Marquette Range Supergroup of the southern Province (Hemlock and Gunflint formations and associated Kiernan sills [#8d]). The 1.88 Ga event or events are of particular importance because they represent a metallotect; the major Ni-Cu-PGE sulphide ores of both the Thompson Nickel Belt and the Raglan deposits of the Cape Smith Belt appear to be associated with this age of magmatism (e.g. Hulbert et al., 2004 for TNB; Wodicka, pers. comm. 2004 for Raglan).

1.88 Ga magmatism in the Cape Smith belt and Labrador Trough has been linked to rifting and possible separation of a microcontinent (St-Onge et al., 2000). However, the magmatism in both the Thompson belt and Animikie basin occurs within a broad

period of ocean closure, and therefore may represent back-arc rifting. The recent discovery of a major ca. 1.88 Ga dyke, the NW-trending Pickle Crow dyke [#8c] crossing the interior of western Superior Province provides a link between the Animikie Basin and Thompson Belt magmatic activity, and may suggest a mantle plume model for the western part of the 1.88 Ga event (Buchan et al., 2003).

1.83–1.82 Ga: The Sparrow dyke swarm [#9a] of the western Rae Province is as yet uncorrelated with any volcanic sequences. Other units of similar age but uncertain relationship are 1) a meta-gabbro and two monzogabbros with U-Pb ages of 1.83–1.82 Ga in the Close Lake, Wollaston-Mudjatik Transition zone [#9a, unit not displayed in Fig. 1], and 2) the widespread Christopher Island formation volcanics of the Baker Basin region [#9a, unit not displayed in Fig. 1].

1.75–1.71 Ga: The Cleaver dykes in the Great Bear Magmatic Zone, the Hadley Bay sills on Victoria Island and MacRae Lake dykes in the northern Rae Province are coeval in age [#10a], suggesting a widespread event in northwestern Canada. This large region of 1.75 Ga activity could be associated with the generation of Wernecke sediments in the northwestern Cordillera, and the approximately 40 Ma younger 1.71 Ga Bonnet Plume River mafic magmatism [#10a; unit not displayed in Fig. 1]. The Wernecke sediments are inferred to represent the earliest rift stage of activity associated with the Cordilleran margin (Cook et al. *in Percival et al.*, 2004). The Pitz formation felsic volcanics and Nueltin felsic intrusive suite [#10a, units not displayed in Fig. 1] of the Baker Basin region are similar in age.

Subhorizontal seismic reflectors interpreted as sills are present over a huge region (about 120,000 sq. km) in the basement underlying the Western Canada Basin. These Winagami sills [#10b] are roughly bracketed in age between 1.89 and 1.76 Ga, and have been tentatively linked with the Cleaver dykes by Ross and Eaton (1997).

1.64 Ga: The 1000 km long Melville Bugt swarm [#11a] is located in western Greenland. Although close to

Baffin Island and other Canadian Arctic islands in a reconstructed Greenland – North America configuration, Melville Bugt dykes have not yet been recognized in Canada.

1.47–1.44 Ga: There are two nodes of activity with this age. The Belt Basin of the western Cordillera contains extensive Moyie-Purcell sills and associated Purcell volcanics [#12a]. These may have been generated when a continent (perhaps Australia) was rifted from the western margin of Laurentia. It is also interesting that in the Wyoming Province there is an extensive linear dyke swarm (Tobacco Root – Group B; [#12a]) of the same age. Assuming that the dykes fed the volcanics and sills, we can infer that the magma source area was either on the western margin of Laurentia in the vicinity of the Belt Basin, or far to the southeast on the southern margin of Laurentia with magma being transported laterally via the dykes to the Belt Basin. Long distance feeding of a sill province has been described above for the 2.22 Ga Ungava event, and may be a not uncommon consequence of the lateral flow pattern in giant dyke swarms (e.g. Ernst and Buchan, 1997).

A second node of this age is located in southeastern Laurentia. The Michael and Shabagamo gabbros [#12b] have an approximate age of 1.47–1.46 Ga and represent a significant event in the southeastern Laurentia. They are located in a back-arc setting to the evolving Grenville Orogen (Rivers et al. *in Percival et al.*, 2004).

1.38 Ga: Hart River volcanic rocks and sills, and coeval sills from the Belt Basin region [#13a], may represent rift sequences (Abbott, 1997; Thorkelson et al., 2003).

1.28–1.27 Ga: One of the largest magmatic events in Canada occurred at 1.267 Ga [#14a]. Most prominently this event consists of the Mackenzie giant radiating dyke swarm. Mackenzie dykes (Fig. 2a) fan over an arc of 100° and cover almost 3 million sq. km of the Canadian Shield. Coeval Coppermine volcanics and the Muskox Intrusion are situated near the plume center, but additional volcanic and sill packages are distributed throughout the swarm and are inferred to be fed via

lateral flow along dykes originating from near the plume center (e.g. Ernst and Baragar, 1992; Baragar et al., 1996). This LIP is presumed to be linked with continental breakup and formation of a northern ocean, which has been termed the Poseidon Ocean (G. Jackson cited in Fahrig, 1987). However, the missing rift block(s) has not yet been identified, although Siberia has been a repeated suggestion (see summary in Ernst et al., 2000; cf. Sears et al., 2004).

The North Atlantic Craton of North America and Greenland is another locus of magmatic activity during this time period [#14b]. The 1.280–1.277 Ga Nain-LP dykes of Labrador are correlative with the BD0 dykes of Greenland (Buchan et al., 1996). Also the 1.273 Ga Harp dykes of Labrador may be linked with Gardar (BD1, BD2, BD3) dykes of Greenland (Baragar, 1977). The setting of this 1.28–1.27 Ga activity is unknown, but it is clearly distinct in location, dyke-trend, and probably in origin from the coeval Mackenzie event. Also widespread in Labrador are the anorthosites, granites, diorites and troctolites of the older and probably unrelated 1.35–1.29 Ga Nain plutonic suite (Ryan and James, 2004 and references therein).

Coeval activity of this age is represented globally by the Central Scandinavian Dolerite Complex (sills) of the Baltic Shield. Paleomagnetic data and geological evidence suggest a reconstructed location east of Greenland (Buchan et al., 2000, and references therein), which is supported by geological evidence (Bingen et al., 2002). Given its great spatial separation from Canada in this reconstruction, the Central Scandinavian Dolerite Complex of Baltica must represent a separate event from the coeval Mackenzie (and also possibly the Nain Province) activity. The occurrence of multiple independent LIPs has been considered evidence for plume cluster events (Ernst and Buchan, 2002).

1.25–1.22 Ga: The Grenville Province and adjacent Superior Province contain widely separated packages of similar-aged magmatism. From southwest to northeast, these include 1.235 Ga Sudbury dykes [#15a], 1.25–1.225 Ga Seal Lake volcanic rocks and

Naskaupi sills [#15b], and the 1.250 Ga Mealy dykes [#15b]. These 1.25–1.22 Ga events along the Grenville Orogen have not been previously linked to each other and more work is required to assess whether they represent disparate elements of the same event. They are situated in a back-arc setting with respect to the evolving Grenville orogen (e.g. Rivers and Corrigan, 2000; Rivers et al. *in* Percival et al., 2004). However a plume origin should not be ruled out. The Seal Lake suite has previously been considered analogous to a flood basalt sequence (“plateau basalt” is the term used in Baragar, 1977), and the aulacogen type geometry of the Sudbury dykes suggests derivation from a spreading center located to the southeast of the swarm (Fahrig, 1987).

1.18–1.14 Ga: Widely separated but possibly linked nodes of mafic magmatism of 1.18 Ga are distributed along the Grenville Orogen. Specifically, toward the northeast end of the Grenville Province the Davy Group sills and dykes [#16a] in the Wakeham Group have an age of 1.177 Ga. Similar ages are found in coronitic gabbros [#16a] in the Baie du Nord segment of Tshenukutish domain. Finally, the same age but with larger uncertainties applies to Algonquin metagabbros [#16a] in the Central Gneiss Belt, southwestern Grenville Province.

Late Gardar magmatism in South Greenland ranges in age from 1.18–1.14 Ga [#16b], and includes the 1.18 and 1.163 Ga Giant Tugtutôq Dykes.

The sparsely distributed 1.141 Ga Abitibi dyke swarm [#16c] of the Central Superior Province has a width of 400 km and extends for nearly 700 km across the southern and eastern Superior Province. This event has long been considered as a precursor to the 1.109–1.085 Ga Keweenaw event [#17a] located at the southwest end of the swarm (next entry). However, the broadly linear pattern of the Abitibi swarm does not eliminate a possible source at the other (northeast end) of the swarm.

1.11–1.08 Ga: One of the most dramatic flood basalt events in Canada and the United States is arguably the Keweenaw magmatism [#17a] of the

Mid-Continent Rift (e.g. Ojakangas et al., 2001). It comprises at least 2×10^6 cubic km of volcanic rocks and possibly an equal volume of intrusive rocks (Fig. 2b). The Mid-Continent Rift (and subsurface lavas) can be traced eastward through Michigan and southwestward into the central United States. A similar age of activity is found in the “Southwestern USA Diabase Province”, and in the Moores Lake sills of the Athabasca Basin (1000 km to the northwest). The Keweenaw activity consists of main pulses of activity at 1.109–1.105 Ga and 1.100–1.094 Ga, but activity is continuous to 1.085 Ga. Emplacement of the Keweenaw LIP is similar in age to that of terminal collision of the Grenville orogen (Rivers et al. *in* Percival et al., 2004). Although a back-arc rifting origin linked to the coeval Grenville orogeny has been suggested, the most widely accepted model links the event to a mantle plume on the basis of the great volume of tholeiitic magma generated in an intraplate setting (e.g. Ojakangas et al., 2001).

0.78 Ga: The Gunbarrel magmatic event [#18a] is distributed over a distance of 2400 km in western North America. Precise 0.780 Ga ages are found in the Hottah sheets of the Slave Province, the Mackenzie Mountains dykes and sills, the MacDonald dykes, and the Tobacco Root- Group B and Wolf Creek sills of the Wyoming Province (Harlan et al., 2003b). The Irene and Huckleberry volcanics of northwestern USA are also inferred to be of this age. The dykes define a radiating swarm with a convergence point in the southern Cordillera near Vancouver Island, indicating a mantle plume origin for this 0.78 Ga LIP (Park et al., 1995). The Windermere sedimentary/volcanic sequence starting at about 0.75 Ga may represent a passive margin associated with this 0.78 Ga plume. Both South China and Australia also contain magmatism dated at 0.78 Ga (events #67 and #64 *in* Ernst and Buchan, 2001, respectively) and both have been proposed as the rifted block(s).

0.72 Ga: Another major LIP event is represented by the Franklin dyke swarm [#19a] which extends throughout the southern Arctic Islands,

but is most significant on Baffin Island, and also on reconstructed Greenland as the Thule swarm [#19a]. The convergence point marking the plume centre for this event is located north or north-west of Banks Island. Natkusiak volcanics, and Minto Inlier and Coronation sills [#19a] are also part of the Franklin event and are generally concentrated toward the plume centre region. The Franklin event may be linked with separation of an as yet unidentified continent.

Another possible locus of similar-aged activity is the Appalachians. A very approximate Rb-Sr age of 0.735 Ga has been suggested for coast-parallel dykes in basement inliers of the Appalachians of the United States (event #72 in Ernst and Buchan, 2001). This activity can be linked with additional magmatism and a stage of rifting along the Laurentian margin in the southern Appalachians (Aleinikoff et al., 1995).

0.62–0.56 Ga (Laurentian margin, Appalachians): The formation of the Iapetus Ocean was preceded by several distinct major magmatic events along the eastern margin of Laurentia. These include the 0.615 Ga Long Range dykes [#20a], the 0.590 Ga Grenville-Adirondack fanning swarm [#20b] and the 0.563 Ga Sept Îles layered intrusion [#20c]. The latter is roughly coeval with the Catocin flood basalts [#20c, unit not displayed in Fig. 1] situated in the southern Appalachians of the United States. Other basaltic units and syenitic intrusions with ages 0.56 to 0.55 Ga are widely distributed (Puffer, 2002; Higgins and van Breemen, 1998).

The oldest event, the 0.615 Ga dykes [#20a] may have been linked to similar-aged magmatism in Baltica, which is represented by the aulacogen-type Egersund dykes (Bingen et al., 1998) and the coast-parallel Baltoscandian breakup swarm (event #55 in Ernst and Buchan, 2001). Two younger rifting events are also recorded in Laurentia: an early separation of Amazonia, and a later separation of the peri-Laurentian Dashwoods microcontinent (Waldron and van Staal, 2001; cf. Cawood et al., 2001). Within this context, the 0.590 Ga radiating swarm [#20b] (associated with the St. Lawrence rift system) may presage the

separation of Amazonia from Laurentia at 0.57 Ga to form the Iapetus Ocean, and the magmatism at 0.563 Ga [#20c] may similarly be linked to the separation of the Dashwoods terrane at 0.550–0.540 Ga.

0.62–0.55 Ga (Avalon zone, Appalachians): The ca. 0.62 Ga Harbour Main [#21a] and the 0.585–0.555 Ga Marystown events [#21b] are recorded in the Avalon zone, which had an uncertain relationship with Laurentia during this period and subsequently drifted with Gondwana before closure of the Iapetus Ocean in the early Paleozoic.

0.57–0.52 Ga (Cordillera): The 0.57 Ga Hamill-Gog Group magmatism [#22a] in the southern Cordillera has been linked with a breakup along this Laurentian margin (Colpron et al., 2002).

In the northern Cordillera lower Paleozoic alkalic and potassic mafic magmatism [#22b] is linked with rifting of the Selwyn Basin (Goodfellow et al., 1995). This magmatism includes the Demster, Menzie Creek, Nidderly volcanics, and younger Fossil Creek volcanics. The only precise age is a 0.518 Ga U-Pb age from a Post-Hyland Group sill, probably representing a widespread set of sills (Abbott, 1997). However, a range of volcanic ages from Lower Cambrian to Early Devonian is suggested on biostratigraphic grounds (Goodfellow et al., 1995; Abbott, 1997).

0.47–0.41 Ga (Dunnage, Gander, Avalon and Meguma zones, Appalachians): In Atlantic Canada several important mafic (and bimodal) magmatic suites include the 0.46 Ga Dunn Point [#23a], and the 0.44–0.43 Ga Bayswater and Cape St. Mary's suites of the Avalon zone [#23b], the Middle Ordovician Overstep Sequence of the Gander and Dunnage zones [#23a], the 0.44 Ga White Rock formation of the Meguma zone [#23b], and the early Silurian Overstep Sequence [#23b] of central Newfoundland (mainly in the Dunnage Zone). Much of this magmatism has been emplaced in a back-arc setting. During this period the Iapetus Ocean was closing, but the spatial relationship between these Iapetan terranes is still under debate.

0.36–0.32 Ga: The Carboniferous Magdalen (or Maritimes) Basin is thought to have been underplated by a layer of mafic magma with an average thickness of 13 km based on geophysical modelling [#24a] (Marillier and Verhoef, 1989). In addition, volcanic rocks and intrusions of this age are distributed widely around the western and southern perimeter of the basin. The Magdalen Basin magmatic event [#24a] has been inferred to represent the final breakthrough of a plume that had been trapped beneath a subducting slab (Murphy et al., 1999).

0.27–0.21 Ga (Cordillera): The mid-Permian to upper Jurassic (ca. 0.27–0.20 Ga) Cache Creek terrane of the Cordillera may include some oceanic plateau and hotspot material [#25a], but this remains controversial (cf. Tardy et al., 2001 and Struik et al., 2001).

Wrangellia is an important accreted terrane of the Cordilleran orogen [#25b], consisting of Karmutsen and Nikolai volcanics and associated 0.232 Ga Maple Creek sills. These units originated as an oceanic plateau possibly formed atop an island arc before being accreted onto the Cordilleran orogen (e.g. Richards et al., 1991).

Note that the Wrangell Lavas (0.065–0.002 Ga) (event #331 in Buchan and Ernst 2004), which extend from the Yukon into Alaska for a distance of ~430 km, are distinct from the much older Wrangellia flood basalts, despite the similar names. The extensive Wrangell lavas are not catalogued in Table 1 because they are linked to subduction along the Aleutian arc.

Ramparts Group magmatism [#25c] of Alaska is poorly dated (ca. 0.21 Ga). It is located in the Tozitna Belt of Alaska and is explained as a "parautochthonous rift assemblage" (p. 190, Dover, 1994).

0.20 Ga: The largest magmatic event on Earth in terms of areal distribution is the Central Atlantic Magmatic Province (CAMP) [#26a]. Found in North America, Europe, Africa and South America, it covers an area of nearly 7 million sq. km. It mainly consists of a giant radiating dyke swarm centred near Florida that was the precursor to the 0.175 Ga opening of

the central Atlantic Ocean. Extensive sill provinces and volcanic packages are also found on all formerly adjacent blocks. In Atlantic Canada, this event is represented by several major dykes, as well as the North Mountain and Grand Manan volcanics [#26a].

A plume origin seems most likely based on the large scale and short duration of the event as well as presence of a giant radiating dyke swarm. However, a model involving "edge convection" has also been advocated mainly on the basis of 'non-plume' chemistry, and the suggestion that the dykes represent a superposition of distinct linear swarms rather than an overall single radiating swarm (e.g. several papers in Hames et al., 2003).

0.14–0.09 Ga: From 0.14 to 0.11 Ga there was extensive NEQ (New England-Quebec) magmatic activity [#27a] in the eastern United States and Quebec. The Monteregian Hills plutons and dykes of this magmatic event have been linked to a plume tail (the Great Meteor hotspot track) associated with the New England seamount chain (Heaman and Kjarsgaard, 2000). Problems with a plume origin for this magmatic province have been discussed by McHone (1996).

Another node of activity at this time is represented by the 400 km long Trap dyke swarm [#27b], which was emplaced along the southwestern coast of Greenland at 0.138 Ga. Given the proximity of Greenland to Labrador at this time, we would expect a continuation of this extensive magmatic event into eastern Canada. Globally, similar-aged magmatism is linked with two plume centres associated with the breakup of South America and Africa (Fig. 8 *in* Ernst and Buchan, 2002).

Magmatism on the Queen Elizabeth Islands [#27c] is part of the High Arctic Large Igneous Province (HALIP) [#27c]. This event is also present in northern Greenland, Svalbard and Franz Josef Land, as well as offshore (Tarduno et al., 1998; Maher, 2001). Precise dates of 0.095–0.092 Ga apply to both the Strand Fiord volcanics and the Wootton intrusion (Tarduno et al., 1998). However K-Ar and Rb-Sr dating as well as biostratigraphic control suggests that the magmatism may have

begun at ca. 0.13 Ga. While a continuum of activity may be present, Maher (2001) interprets two pulses of activity, at ca. 0.13 and ca. 0.09 Ga. The identification of a radiating dyke swarm suggests a plume origin with a plume centre located near the Alpha Ridge (Embry and Osadetz, 1988; Ernst and Buchan, 1997; Maher, 2001).

0.07–0.05 Ga: The 0.070 Ga Carmacks volcanics [#28a] of the northern Cordillera have a slightly more potassic composition than most of the other events discussed in this paper, but they have a large areal extent (about 60,000 sq. km) and were apparently emplaced in a short duration. The slightly younger 0.06–0.05 Ga Crescent Terrane volcanics [#28b] consist of thick basaltic sequences distributed over a ca. 600 km distance along the Coast Ranges of western North America. They are inferred to represent accreted seamounts. Both the Carmacks volcanics [#28a] as well as the Crescent Terrane volcanics [#28b] have been interpreted as originating from earlier stages of the Yellowstone plume, prior to its main expression as the Columbia River LIP at 0.017 Ga (see next entry, [#29b]) (e.g. Johnston et al., 1996; Murphy et al., 2003).

The North Atlantic Igneous Province (NAIP) [#28c] is most voluminous in eastern Greenland and the UK, and adjacent offshore regions. Minor activity at Cape Dyer and Cape Searle on Baffin Island, as well as more substantial magmatism in west Greenland are linked with the NAIP. The NAIP LIP is linked to the present-day Icelandic hotspot.

0.025–0.015 Ga: An important intraplate event is the widespread Behm Canal event [#29a] of the Cordillera, which is also known as the Tertiary Lamprophyre Province. It consists of alkaline lamprophyres that are considered volatile-enriched equivalents of alkali basalts (Rock, 1991, p. 11-12, 122).

The voluminous Columbia River Basalt Group (CRBG) LIP [#29b] of the northwestern United States was mainly erupted from 0.017 to 0.015 Ga. It has been linked to the Yellowstone plume. The coeval Chilcotin Group volcanics [#29c] are located a short distance to

the north in Canada, and although they are areally extensive they are volumetrically minor.

Archean

Archean greenstone belts represent deformed and fragmented volcanic suites, and are of two main affinities: calc-alkaline and tholeiitic (e.g. de Wit and Ashwal, 1997; Condie, 2001; Bleeker, 2002). The calc-alkaline suites are considered to be of arc origin whereas the tholeiitic suites, particularly those containing komatiites, are not. The presence of komatiites satisfies one requirement for the identification of LIPs, i.e. that they not be produced by subduction. In addition, greenstone belts with komatiites are probably not produced by normal spreading ridge processes because komatiites indicate source region temperatures higher than those associated with normal spreading ridges. There is some controversy on this point since the Archean geotherm was hotter. The scale of Archean LIP candidates is also uncertain. In most cases deformation and faulting prevents the recognition of Archean tholeiite-komatiite greenstone belts over LIP-scale distances. So the Archean LIP history, discussed below and summarized in Table 2 and Figure 1, remains speculative. Events are included in Table 2 on the basis of either the presence of komatiites, and/or inferred oceanic plateau setting.

3.11–2.98 Ga: The oldest known period of potential LIP activity is represented by komatiite-bearing greenstone belts (3.105 Ga Hunt River [#A1a] and 2.99–2.98 Ga Florence Lake [#A1b] belts) located in the Hopedale block of the Nain Province of the North Atlantic Craton. Another important magmatic event is associated with the ca. 2.99 Ga rifting [#A1c] of the ancient Archean nucleus in the western Superior Province comprising the North Caribou, Central Wabigoon and Marmion blocks (Tomlinson et al., 1999; Tomlinson and Condie, 2001).

2.93–2.92 Ga: A subsequent stage of komatiite-bearing greenstone belts [#A2a] at about 2.93–2.92 Ga is also widespread in the western Superior Province, but the setting and link with associated arc-type greenstone belts is unclear.

2.86 Ga: The Pickle Crow greenstone belt [#A3a] in the Uchi subprovince is approximately dated at 2.86 Ga and contains komatiites.

2.79–2.77 Ga: The 2.786 Ga Vizion greenstone belt [#A4a] of northern Quebec contains komatiites and has been linked with formation of an oceanic plateau. Greenstone belt fragments (containing komatiites) within the Faribault-Thury Complex [#A4b] are also situated in northern Quebec, and are bracketed in age between 2.785 and 2.710 Ga. (Note that if future dating determines an age closer to the younger end of the age bracket, then the Faribault-Thury Complex event would be grouped in the next entry.)

The 2.775 Ga Fourbay sequence [#A4c] of the western Superior Province lacks komatiites, but is included in this LIP compilation because of its inferred oceanic plateau origin.

2.75–2.70 Ga: In various portions of the Canadian Shield there was widespread mafic magmatism falling in the age range 2.75–2.70 Ga. This is discussed in terms of three regional groupings.

It has been suggested that the Prince Albert Group [#A5a] can be correlated with the Woodburn and Mary River groups, thus defining a single event in the Rae Province of northern Canada, which is distributed over a lateral distance of nearly 1000 km. The magmatism which includes komatiites is interpreted to have initiated at 2.73 Ga and may be associated with plume-generated continental breakup.

The Kam Group [#A5b] has been traced across large parts of the Slave Province through a complicated deformation pattern (Bleeker, 2003, and references therein). The magmatism of the Kam Group ranges from 2.734–2.700 Ga, and was particularly voluminous from 2.72–2.70 Ga.

Magmatism in the Abitibi greenstone belt [#A5c] of the central Superior Province is widespread and consists of four distinct stages of komatiite-associated magmatism emplaced during the interval 2.75–2.70 Ga (e.g. Sproule et al., 2002). The interspersed calc-alkaline magmatism

suggests a plume arc association (e.g. Wyman, 1999). The Schreiber-Hemlo-White River-Dayohessarah packages found in the Wawa greenstone belt [#A5d], also contain significant komatiite-bearing tholeiitic magmatism with ages between 2.75 and 2.73 Ga. The Wawa belt is on strike with and is probably the continuation of the Abitibi belt.

DISCUSSION

As described above, Canada has a rich LIP history consisting of at least 80 possible events (Fig. 1, Tables 1 and 2). They range from Archean greenstone belts (containing komatiites) such as the Prince Albert Group and probable correlatives, which may extend for >1000 km along the Rae Province, through Proterozoic giant radiating dyke swarms, such as the Mackenzie swarm that covers nearly 3,000,000 sq. km of the Canadian Shield, to young flood basalts, such as the rift-related Keweenaw Group and the accreted oceanic plateau, Wrangellia, in the Cordillera. LIPs are key to resolving a number of important geological issues and processes. Here we apply our database of Canadian LIPs to several frontier issues.

Plume vs. Non-plume Origins

As mentioned earlier, there is currently an intense debate about plume versus non-plume origins for LIPs. This debate is occurring both on the web (e.g. www.mantleplumes.org; www.largeigneousprovinces.org) and in the scientific literature (e.g. Anderson, 2001; Foulger and Natland, 2003; DePaolo and Manga, 2003; Ernst et al., 2004). The Canadian LIP database can contribute to this debate in various ways. Giant radiating dyke swarms are strongly indicative of mantle plumes (e.g. Ernst and Buchan, 1997). Using this criteria, plumes would be inferred at 2.45, 1.27, 0.78, 0.72, 0.20 Ga. Furthermore, many Canadian swarms have a failed-arm (aulacogen setting) (Fig. 3b, Table 3), which also suggests plume involvement. In addition, Archean greenstone belts, which contain komatiites, are arguably plume-related. Finally, recognition of additional mantle plumes may also derive from studies of

regional uplift patterns (see below). On the other hand, some LIPs having a linear distribution, such as those in a back-arc setting, may be consistent with non-plume origins. Those along a breakup margin may be generated by decompression melting accompanying rifting. Some LIPs may consist of two pulses, an initial burst of magmatism associated with plume arrival and a second caused by the onset of decompression melting associated with breakup.

Precise Time Markers for Stratigraphic Correlation

The wide distribution (potentially over millions of sq. km) and the typically short duration of events makes them ideal as precise stratigraphic markers (e.g. LeCheminant and Heaman, 1989; Harlan et al., 2003b). For instance, recognition of the same magmatic event within widely separated sedimentary sequences represents an ideal marker for inter-basin correlation. In Canada, the 1.27 Ga Mackenzie, 0.78 Ga Gunbarrel, and 0.72 Ga Franklin LIPs represent particularly good markers.

Reconstruction of Continents

Globally there is a clear link between young LIPs and breakup margins (e.g. Courtillot et al., 1999). The Canadian landmass preserves a history of continental breakup, and subsequent reassembly marked by sutures. Therefore, the Canadian LIP record is fertile ground for exploring links with breakup events. Archean continental fragments each contain a particular age distribution of mafic events (mostly Proterozoic dyke swarms) that represent a distinct “bar code” (Bleeker, 2003, 2004). Comparison of the “bar code” from the approximately 35 different Archean continental fragments (at least eight of which are in Canada) represents a key tool for proposing reconstructions between these fragments. Reconstructions can be tested by comparing the paleomagnetism of coeval mafic units on the different continental fragments (Buchan et al., 2000). In addition, linear (Fig. 4a) and radiating (Fig. 4b) dyke swarms can be used to constrain the reconstruction geometry. Some suggested correlations are given in




Figure 4 Use of giant dyke swarms for continental reconstruction **a.** Giant linear dyke swarms used as piercing points. **b.** Giant radiating swarms.

Tables 1 and 2, and in the accompanying text.

Exploration (Ni-Cu-PGEs and Diamonds)

Ni-Cu-PGE deposits are commonly associated with LIPs. Notable examples include the Siberian Traps and the Bushveld Complex (e.g. Naldrett, 1999; Pirajno, 2000; Diakov et al., 2002; Hulbert, 2002). Ernst and Hulbert (2003) carried out a preliminary analysis of background PGE levels in about 60 Canadian LIPs and other intraplate mafic events in order to assess which are more likely to host such deposits. Events with elevated background PGE levels (≥ 10 ppb Pt and Pd) are thought to have greater potential for enrichment during magmatic emplacement. Ernst and Hulbert (2003) found that events with high background levels include the 2.50–2.45 Ga Matachewan, 2.22–2.21 Ga Ungava-Nipissing, 1.27 Ga Mackenzie, 0.72 Ga Franklin, 0.59 Ga Grenville and portions of the 1.11–1.09 Ga Keweenawan and 0.13–0.09 Ga Sverdrup Basin events.

Some studies have proposed a direct link between kimberlites and underlying plumes (Haggerty, 1999;

Heaman and Kjarsgaard, 2000; Schissel and Smal, 2001). Recently it has also been suggested that some kimberlites might be preferentially localized along particular dyke swarms such as the 2.023 Ga Lac de Gras swarm of the Slave Province (Wilkinson et al., 2001; Stubble, 2003) and the 2.50–2.45 Ga Matachewan dykes of the Attawapiskat region in the James Bay lowlands (Stott and Halls, 2002). Further work is required to test the potential of fractures and zones of weakness represented by major dyke swarms to localize kimberlite magmas during their ascent through the crust.

Climate Change

Numerous studies have explored the link between LIPs and climate change (e.g. Condie, 2001; Isley and Abbott, 2002; Courtillot and Renne, 2003; Ernst and Buchan, 2003; Prokoph et al., 2003). The correlation between extinction events and LIPs is compelling (Courtillot et al., 1996; Courtillot and Renne, 2003) and suggests that some LIPs, perhaps acting in concert with meteorite impact events, may be the trigger for extinction events. Evaluation of a robust Canadian LIP record can contribute to the global understanding of the effect of LIPs on climate. For instance, all other factors being equal, the largest LIP events should have the largest climatic effect. Some of the largest LIPs in Canada are the ca. 1.88 Ga Circum-Superior, 1.27 Ga Mackenzie, 0.72 Ga Franklin, 0.615–0.555 Ga Central Iapetus, and 0.20 Ga CAMP events. Only the climatic effect of the youngest of these events has been evaluated (Pálffy 2003).

Regional Domal Uplift

LIPs linked to mantle plumes should be associated with regional domal uplift. The scale of uplift corresponds to the size of the plume head (e.g. Cox, 1989; Rainbird and Ernst, 2001; Campbell, 2001; Şengör, 2001; He et al., 2003). The largest plumes are thought to generate uplifts with a radius 1000 km and a peak elevation 1 to 2 km. Such uplifts should exert a first-order control on concurrent regional sedimentation patterns. Plume head related uplift has only received preliminary investigation

in Canada, where it has been associated with 1.27 Ga Mackenzie, 0.72 Ga Franklin, and 0.615–0.555 Ga Central Iapetus events (e.g. Rainbird and Ernst, 2001). It is hoped that the LIP database for Canada will stimulate further investigation of regional uplift patterns.

EXPANDING THE LIP DATABASE TO ADDRESS FRONTIER ISSUES

The frontier issues discussed above can only be fully addressed using a more robust LIP record. Although an extensive database is presented in this paper, many major mafic and ultramafic units remain undated and poorly characterized. Rapid improvement in the LIP database can be achieved only through a concerted campaign of geochronology integrated with other fields such as paleomagnetism and geochemistry, as is being proposed for Canada by Bleeker (2004) and internationally by Ernst et al. (2004).

ACKNOWLEDGEMENTS

John Morgan helped compile information on the link between Canadian LIPs and breakup events. Discussions with Tom Skulski and Wouter Bleeker have been invaluable in guiding our application of LIP concepts to Archean greenstone belts. Mike Hamilton, Tony LeCheminant and Henry Halls are acknowledged for numerous discussions on the nature and setting of Proterozoic dyke swarms of Canada. Thoughtful reviews were provided by Wouter Bleeker, Michael Higgins and Peter Lightfoot. Figure 2a (portion of aerial photograph #A14233-138) © 2004, Her Majesty the Queen in Right of Canada, reproduced from the collection of the National Air Photo Library with permission of Natural Resources Canada. Figure 2b (archive number 202781) is reproduced with the permission of the Minister of Public Works and Government Services Canada, 2004, and courtesy of Natural Resources Canada, Geological Survey of Canada.

REFERENCES

- Aleinikoff, J.N., Zartman, R.E., Walter, M., Rankin, D.W., Lyttle, P.T. and Burton, W.C., 1995, U-Pb ages of metarhyolites of the Catocin and Mount Rogers formations,

- central and southern Appalachians: Evidence for two pulses of Iapetan rifting: *American Journal of Science*, v. 295, p. 428-454.
- Anderson, D.L., 2001, Top-down tectonics: *Science*, v. 293, p. 2016-2018.
- Anderson, R.G., Resnick, J., Russell, J.K., Woodsworth, G.J., Villeneuve, M.E. and Grainger, N.C., 2001, The Cheslatta Lake suite: Miocene mafic, alkaline magmatism in central British Columbia: *Canadian Journal of Earth Sciences*, v. 38, p. 697-717.
- Annesley, I.R., Madore, C., Kamo, S.L., Kwok, Y.Y., Cutts, C. and Portella, P., 2003, U-Pb zircon and monazite geochronology of tonalitic gneiss from Close Lake, Wollaston-Mudjatik Transition Zone, northern Saskatchewan [abstract], Vancouver 2003 (GAC-MAC-SEG meeting), abstract no. 545 CD-ROM.
- Arndt, N.T., Ginibre, C., Chauvel, C., Albarède, F., Cheadle, M., Herzberg, C., Jenner, G., and Lahaye, Y., 1998, Were komatiites wet?: *Geology*, v. 26, p. 739-742.
- Aspler, L.B., Chiarenzelli, J. R., Cousens, B.L. and Valentino, D., 1999, Precambrian geology, northern Angikuni Lake, and a transect across the Snowbird tectonic zone, western Angikuni Lake, Northwest Territories (Nunavut): *in* Current Research, Part C, Geological Survey of Canada Paper 1999-C, p. 107-118.
- Aspler, L.B., Cousens, B.L. and Chiarenzelli, J.R., 2002, Long-distance intracratonic transport of mafic magmas during opening of the Manikewan ocean (Trans-Hudson orogen): Griffin gabbro sills (2.11 Ga), Hurwitz Basin, Nunavut, Canada: *Precambrian Research*, v. 117, p. 269-294.
- Ayer J., Amelin, Y., Corfu, F., Kamo, S., Ketchum, J. et al., 2002, Evolution of the southern Abitibi greenstone belt based on U-Pb geochronology: autochthonous volcanic construction followed by plutonism, regional deformation and sedimentation: *Precambrian Research*, v. 115, p. 63-95.
- Baragar, W.R.A., 1977, Volcanism of the stable crust, *in* Baragar, W.R.A., Coleman, L.C. and Hall, J.M. eds., *Volcanic Regimes in Canada: Geological Association of Canada, Special Paper 16*, p. 377-405.
- Baragar, W.R.A. and Scoates, R.F.J., 1987, Volcanic geochemistry of the northern segments of the Circum-Superior Belt of the Canadian Shield, *in* Pharaoh, T.C., Beckinsale, R.D and Rickard, D. (eds.), *Geochemistry and Mineralization of Proterozoic Volcanic Suites: Geological Society Special Publication* (London), no. 33, p. 113-131.
- Baragar, W.R.A, Ernst, R.E., Hulbert, L. and Peterson, T., 1996, Longitudinal petrochemical variation in the Mackenzie dyke swarm, northwestern Canadian Shield: *Journal of Petrology*, v. 37, p. 317-359.
- Barr, S.M., White, C.E. and McLeod, M.J., 1999, Geology of the Silurian Kingston terrane, southern New Brunswick; *in* Carroll, B.M.W. eds., *Current Research 1998: New Brunswick Department of Natural Resources and Energy, Minerals and Energy Division, Mineral Resources Report 99-4*, p. 1-17.
- Bell, R., 1870, Report of Mr. Robert Bell on Lakes Superior and Nipigon: Geological Survey of Canada, Report of Progress 1866-69, Part IX, p. 313-364.
- Bell, R., 1879, Report on an exploration of the east coast of Hudson's Bay: Geological Survey of Canada, Report of Progress 1877-78, Part C, p. 1-37.
- Bevier, M.L., Barr, S.M., White, C.E. and Macdonald, A.S., 1993, U-Pb geochronologic constraints on the volcanic evolution of the Mira (Avalon) terrane, southeastern Cape Breton Island, Nova Scotia: *Canadian Journal of Earth Sciences*, v. 30, p. 1-10.
- Bingen, B., Demaiffe, D. and van Breemen, O., 1998, The 616 Ma old Egersund basaltic dike swarm, SW Norway, and Late Neoproterozoic opening of the Iapetus Ocean: *Journal of Geology*, v. 106, p. 565-574.
- Bingen, B., Mansfeld, J., Sigmond, E.M.O. and Stein, H., 2002, Baltica-Laurentia link during the Mesoproterozoic: 1.27 Ga development of continental basins in the Sveconorwegian Orogen, southern Norway: *Canadian Journal of Earth Sciences*, v. 39, p.1425-1440.
- Bleeker, W., 2002, Archaean tectonics: a review with illustrations from the Slave craton, *in* Fowler, C.M.R., Ebinger, C.J., and Hawkesworth, C.J., (eds.), *The Early Earth: Physical, Chemical and Biological Development: Geological Society, London, Special Publications* v. 199, p. 151-181.
- Bleeker, W., 2003, The later Archean record: a puzzle in ca. 35 pieces: *Lithos*, v. 970, p. 99-134.
- Bleeker, W., 2004, Taking the pulse of planet Earth: a proposal for a new multi-disciplinary flagship project in Canadian solid Earth sciences: *Geoscience Canada* (in press)
- Bleeker, W. and Kamo, S., 2003, A precise age for the Duck Lake sill and its relevance for fitting the Slave in a global Archean context: *in* 31st Yellowknife Geoscience Forum, November 19-21, C.S. Lord Northern Geoscience Centre, Yellowknife, Canada.
- Buchan, K.L. and Ernst, R.E., 2004, Diabase dyke swarms and related units in Canada and adjacent regions: Geological Survey of Canada, Map 2022A (1:5,000,000 map with accompanying compilation).
- Buchan, K.L. and Halls, H.C., 1990, Paleomagnetism of Proterozoic mafic dyke swarms of the Canadian Shield, *in* Parker, A.J., Rickwood, P.C. and Tucker, D.H. (eds.), *Mafic Dykes and Emplacement Mechanisms: Balkema, Rotterdam*. p. 209-230.
- Buchan, K.L., Harris, B.A., Ernst, R.E. and Hanes, J.A., 2003, Ar-Ar dating of the Pickle Crow diabase dyke in the western Superior craton of the Canadian Shield of Ontario and implications for a possible plume centre associated with ca. 1880 Ma Molson magmatism of Manitoba: *Geological Association of Canada / Mineralogical Association of Canada Abstracts*. (no. 481 on CD-ROM, p. 17 in print volume).
- Buchan, K.L., Hodych, J.P., Roddick, J.C., Emslie, R.F. and Hamilton, M.A., 1996, Paleomagnetism and U-Pb geochronology of Mesoproterozoic dykes of Labrador and correlations with dykes of southwest Greenland, *in* Proterozoic Evolution of the North Atlantic Realm (Program and Abstracts): conference held in Goose Bay, Labrador, Canada and organized by IGCP 371, ECSOOT Lithoprobe Canada and International Basement Tectonics Association, p. 37.
- Buchan, K.L., Goutier, J., Hamilton, M.A., Ernst, R.E. and Matthews, W., 2004, Paleomagnetism and U-Pb geochronology of Lac Esprit dykes of Quebec and implications for Paleoproterozoic deformation of the Superior Province: Abstract for joint AGU-CGU annual meeting, Montreal.
- Buchan, K.L., Mortensen, J.K., Card, K.D. and Percival, J.A., 1998, Paleomagnetism and U-Pb geochronology of diabase dyke swarms of Minto block, Superior Province, Quebec, Canada: *Canadian Journal of Earth Sciences*, v. 35, p. 1054-1069.
- Buchan, K.L., Mertanen, S., Park, R.G., Pesonen, L.J., Elming, S.Å., Abrahamsen, N. and Bylund, G., 2000, Comparing the drift of Laurentia and Baltica in the Proterozoic: The importance of key paleomagnetic poles: *Tectonophysics*, v. 319, p. 167-198.
- Campbell, I.H., 1998, The mantle's chemical structure: insights from the melting products of mantle plumes, *in* Jackson, I.N.S. (ed.), *The Earth's mantle: Composition, structure and evolution*. Cambridge University Press, Cambridge, p. 259-310.
- Campbell, I.H., 2001, Identification of ancient mantle plumes, *in* Ernst, R.E. and Buchan, K.L., eds. *Mantle Plumes: Their Identification Through Time: Geological Society of America, Special Paper 352*, p. 5-21.
- Campbell, I.H. and Griffiths, R.W., 1990, Implications of mantle plume structure for the evolution of flood basalts: *Earth and Planetary Science Letters*, v. 99, p. 79-93.
- Cawood, P.A., McCausland, P.J.A. and Dunning, G.R., 2001, Opening Iapetus: constraints from the Laurentian margin in Newfoundland: *Geological Society of America, Bulletin*, v. 113, p. 443-453.
- Chandler, F.W. and Schwarz, E.J., 1980, Tectonics of the Richmond Gulf area, northern Quebec – A hypothesis, *in* Current Research, Part C; Geological Survey of Canada, Paper 80-1C, p. 59-68.
- Coffin M.F. and Eldholm, O., 1994, Large igneous provinces: crustal structure, dimensions, and external consequences:

- Review of Geophysics, v. 32, p. 1-36.
- Coffin M.F. and Eldholm, O., 2001, Large igneous provinces: progenitors of some ophiolites?, *in* Ernst, R.E. and Buchan, K.L., (eds.), *Mantle Plumes: Their Identification Through Time*: Geological Society of America, Special Paper 352, p. 59-70.
- Colpron, M., Logan, J.M. and Mortensen, J.K., 2002, U-Pb zircon age constraint for late Neoproterozoic rifting and initiation of the lower Paleozoic passive margin of western Laurentia: *Canadian Journal of Earth Sciences*, v. 39, p. 133-143.
- Condie, K.C., 2001, *Mantle Plumes and their Record in Earth History*: Cambridge University Press, Oxford U.K. 306 p.
- Corrigan, D., Rivers, T. and Dunning, G., 2000, U-Pb constraints for the plutonic and tectonometamorphic evolution of Lake Melville terrane, Labrador and implications for basement reworking in the northeastern Grenville Province: *Precambrian Research*, v. 99, p. 65-90.
- Courtillot, V.E. and Renne, P.R., 2003, On the ages of flood basalt events: *C.R. Geoscience*, v. 335, p. 113-140.
- Courtillot, V., Davaille, A., Besse, J. and Stock, J., 2003, Three distinct types of hotspots in the Earth's mantle: *Earth and Planetary Science Letters*, v. 205, p. 295-308.
- Courtillot, V., Jaeger, J.J., Yang, Z., Féraud, G. and Hofmann, C., 1996, The influence of continental flood basalts on mass extinctions: where do we stand? *in* Ryder, G., Fastovsky, D. and Gartner, S. (eds.), *The Cretaceous-Tertiary Event and Other Catastrophes in Earth History*: Geological Society of America, Special Paper 307, Boulder CO, p. 513-525.
- Courtillot, V., Jaupart, C., Manighetti, I., Tapponnier, P. and Besse, J., 1999, On causal links between flood basalts and continental breakup: *Earth and Planetary Science Letters*, v. 166, p. 177-195.
- Cousens, B.L., Aspler, L.B., Chiarenzelli, J.R., Donaldson, J.A., Sandeman, H., Peterson, T.D., LeCheminant, A.N., 2001, Enriched Archean lithospheric mantle beneath western Churchill Province tapped during Paleoproterozoic orogenesis: *Geology*, v. 29, p. 827-830.
- Cox, K.G., 1989, The role of mantle plumes in the development of continental drainage patterns: *Nature*, v. 342, p. 873-877.
- DePaolo, D. and Manga, M., 2003, Deep origin of hotspots – Is only seeing believing?: *Science*, v. 300, p. 920-921.
- de Wit, M.J., and Ashwal, L.D., 1997, *Greenstone belts*: Oxford, UK, Clarendon 809 p.
- Dover, J.H., 1994, *Geology of part of east-central Alaska*, *in* Plafker, G. and Berg, H.C. (eds.), *The Geology of North America*: v. G-1, p. 153-204.
- Diakov, S., West, R., Schissel, D., Krivtsov, A., Kochnev-Pervoukhov, V. and Migachev, I., 2002, Recent advances in the Noril'sk model and its application for exploration of Ni-Cu-PGE sulfide deposits, *in* Goldfarb, R.J. and Nielsen, R.L. (eds.), *Integrated Methods for Discovery: Global Exploration in the Twenty-First Century*: Society of Economic Geologists, Special Publication 9, p. 203-226.
- Embry, A.F. and Osadetz, K.G., 1988, Stratigraphy and tectonic significance of Cretaceous volcanism in Queen Elizabeth Islands, Canadian Arctic Archipelago: *Canadian Journal of Earth Sciences*, v. 25, p. 1209-1219.
- Eriksson, P.G., Condie, K.C., van der Westhuizen, W., van der Merwe, R., de Bruijn, H., et al., 2002, Late Archaean superplume events: a Kaapvaal-Pilbara perspective: *Journal of Geodynamics*, v. 34, p. 207-247.
- Ernst, R.E. and Baragar, W.R.A., 1992, Evidence from magnetic fabric for the flow pattern of magma in the Mackenzie giant radiating dyke swarm: *Nature*, v. 356, p. 511-513.
- Ernst, R.E. and Buchan, K.L., 1997, Giant radiating dyke swarms: their use in identifying pre-Mesozoic large igneous provinces and mantle plumes, *in* Mahoney, J.J. and Coffin, M.F., (eds.), *Large Igneous Provinces: Continental, Oceanic, and Planetary Flood Volcanism*: American Geophysical Union, Geophysical Monograph 100, p. 297-333.
- Ernst, R.E. and Buchan, K.L., 2001, Large mafic magmatic events through time and links to mantle plume heads, *in* Ernst, R.E. and Buchan, K.L. (eds.), *Mantle Plumes: Their Identification Through Time*: Geological Society of America, Special Paper 352, p. 483-575.
- Ernst, R.E. and Buchan, K.L., 2002, Maximum size and distribution in time and space of mantle plumes: evidence from large igneous provinces, *in* Condie, K.C., Abbot, D. and Des Marais, D.J. (eds.), *Superplume Events in Earth's History: Causes and Effects*: *Journal of Geodynamics (Special Issue)*, v. 34, p. 309-342 [Erratum, *J. Geodynamics* 2002, v. 34, p. 711-714].
- Ernst, R.E. and Buchan, K.L., 2003, Recognizing mantle plumes in the geological record: *Annual Review of Earth and Planetary Sciences*, v. 31, p. 469-523.
- Ernst, R.E. and Hulbert, L.J., 2003, Background Pt-Pd levels in mafic Large Igneous Provinces (LIPs) in Canada: *Geological Survey of Canada, Open File 4472 (Poster)*.
- Ernst, R.E., Buchan, K.L., Hamilton, M.A., Okrugin, A.V. and Tomshin, M.D., 2000, Integrated paleomagnetism and U-Pb geochronology of mafic dykes of eastern Anabar Shield region, Siberia: implications for Mesoproterozoic paleolatitude of Siberia and comparisons with Laurentia: *Journal of Geology*, v. 108, p. 381-401.
- Ernst, R.E., Buchan, K.L. and Campbell, I.H., 2004, *Frontiers in large igneous province (LIP) research*: Lithos (in press).
- Fahrig, W.F., 1987, The tectonic settings of continental mafic dyke swarms: failed arm and early passive margin, *in* Halls, H.C. and Fahrig, W.F., eds., *Mafic Dyke Swarms*: Geological Association of Canada Special Paper 34, p. 331-348.
- Ferri, F., Rees, C.J., Nelson, J.L., and Legun, A.S., 1999, Geology and mineral deposits of the northern Kechika Trough between Gataga River and the 60th parallel: *British Columbia Ministry of Energy and Mines, Bulletin* 107, 122 p.
- Foulger, G.R., and Natland, J.H., 2003, Is "hotspot" volcanism a consequence of plate tectonics?: *Science*, v. 300, p. 921-922.
- Fralick, P., Davis, D.W. and Kissin, S.A., 2002, The age of the Gunflint Formation, Ontario, Canada: single zircon U-Pb age determination from reworked volcanic ash: *Canadian Journal of Earth Sciences*, v. 39, p. 1085-1091.
- French, J.E., Heaman, L.M., and Chacko, T., 2002, Feasibility of chemical U-Th-total Pb baddeleyite dating by electron microprobe: *Chemical Geology*, v. 188, p. 85-104.
- Goodfellow, W.D., Cecile, M.P. and Leybourne, M.L., 1995, Geochemistry, petrogenesis, and tectonic setting of lower Paleozoic alkalic and potassic volcanic rocks, Northern Canadian Cordilleran Miogeocline: *Canadian Journal of Earth Sciences*, v. 32, p. 1236-1254 [Erratum: v. 32, p. 2167].
- Gower, C.F. and Krogh, T.E., 2002, A U-Pb geochronological review of the Proterozoic history of the eastern Grenville Province: *Canadian Journal of Earth Sciences*, v. 39, p. 795-829.
- Haggerty, S.E., 1999, A diamond trilogy: superplumes, supercontinents, and supervolcanoes: *Science*, v. 285, p. 851-860.
- Halls, H.C., 1982, The importance and potential of mafic dyke swarms in studies of geodynamic processes: *Geoscience Canada*, v. 9, p. 145-154.
- Halls, H.C. and Davis, D.W., 2004, Paleomagnetism and U-Pb geochronology of the 2.17 Ga Biscotasing dyke swarm, Ontario, Canada: Evidence of vertical-axis crustal rotation across the Kapuskasing Zone: *Canadian Journal of Earth Sciences*, v. 41, p. 255-269.
- Halls, H.C. and Heaman, L.M., 2000, The paleomagnetic significance of new U-Pb age data from the Molson dyke swarm, Cauchon Lake area, Manitoba: *Canadian Journal of Earth Sciences*, v. 37, p. 957-966.
- Hames, W.E., McHone, J.G., Renne, P.R. and Ruppel, C. (eds.), 2003, *The Central Magmatic Province: Insights from Fragments of Pangea*: American Geophysical Union Geophysical Monograph Series 136, 267 p.
- Hamilton, M.A., and Murphy, J.B., 2004, Tectonic significance of a Llanvirn age for the Dunn Point volcanic rocks, Avalon

- terrane, Nova Scotia, Canada: implications for the evolution of the Iapetus and Rheic oceans: *Tectonophysics*, v. 379, p. 199-209.
- Hamilton, M.A., Buchan, K.L., Garde, A.A. and Connelly, J.N., 2004, U-Pb age and preliminary paleomagnetism of a Melville Bugt diabase dyke, West Greenland, and implications for mid-Proterozoic Laurentia-Baltica reconstructions: Abstract for the joint AGU-CGU annual meeting, Montreal.
- Hanson, R.E., Crowley, J.L., Bowring, S.A., Ramezani, J., Goes, W.A., Dalziel, I.W.D., Pancake J.A., Seidel, E.K., Blenkinsop, T.G. and Mukwakwami, J., 2004, Coeval large-scale magmatism in the Kalahari and Laurentian cratons during Rodinia assembly: *Science*, 304, p. 1126-1129.
- Harlan, S.S., Geissman, J.W. and Premo, W.R., 2003a, Paleomagnetism and geochronology of an Early Proterozoic quartz diorite in the southern Wind River Range, Wyoming, USA: *Tectonophysics*, v. 362, p. 105-122.
- Harlan, S.S., Heaman, L., LeCheminant, A.N. and Premo, W.R., 2003b, Gunbarrel mafic magmatic event: A key 780 Ma time marker for Rodinia plate reconstructions: *Geology*, v. 31, p. 1053-1056.
- Hartlaub, R.P., Ashton, K.E., Heaman, L.M. and Chacko, T., 2002, Was there an ~2000 km long Neoproterozoic extensional event in the Rae Craton? Evidence from the Murmac Bay Group of northern Saskatchewan: [abstract]: Saskatoon 2002, Joint annual meeting of the Geological Association of Canada, and Mineralogical Association of Canada, Saskatoon Saskatchewan, Canada, May 27-29, 2002.
- He, B., Xu, Y.-G., Chung, S.-L., Xiao, L., and Wang, Y., 2003, Sedimentary evidence for a rapid crustal doming prior to the eruption of the Emeishan flood basalts: *Earth and Planetary Science Letters*, v. 213, p. 389-403.
- Heaman, L.M., 1997, Global mafic magmatism at 2.45 Ga: Remnants of an ancient large igneous province?: *Geology*, v. 25, p. 299-302.
- Heaman, L.M., and Kjarsgaard, B.A., 2000, Timing of eastern North American kimberlite magmatism: continental extension of the Great Meteor hotspot track?: *Earth and Planetary Science Letters*, v. 178, p. 253-268.
- Heaman, L.M., Erdmer, P., and Owen, J.V., 2002, U-Pb geochronologic constraints on the crustal evolution of the Long Range Inlier, Newfoundland: *Canadian Journal of Earth Sciences*, v. 39, p. 845-865.
- Higgins, M.D. and van Breemen, O., 1998, The age of the Sept Iles layered mafic intrusion, Canada: implications for the Late Neoproterozoic/Cambrian history of southeastern Canada: *The Journal of Geology*, v. 106, p. 421-431.
- Hodych, J.P., and Buchan, K.L., 1998, Palaeomagnetism of the ca. 440 Ma Cape St. Mary's sills of the Avalon Peninsula of Newfoundland: implications for Iapetus Ocean closure: *Geophysical Journal International*, v. 135, p. 155-164.
- Hollings, P.N., Wyman, D. and Kerrich, R., 1999, Komatiite-basalt-rhyolite volcanic association in northern Superior Province greenstone belts: significance of plume-arc interaction in the generation of the proto continental Superior Province: *Lithos*, v. 46, p. 137-161.
- Hulbert, L., 2002, Magmatic platinum group element environments in Canada: present and future exploration target areas: Geological Association of Canada Robinson Lecture 2002, MP#2 [CD-ROM]
- Hulbert, L., Stern, R., Kyser, T.K., Pearson, J., Leshner, M. and Grinenko, L., 1994, The Winnipegosis komatiite belt Central Manitoba. Manitoba Energy and Mines, Manitoba: Mining Minerals and Petroleum Convention 94. Program with Abstracts, p. 21.
- Hulbert, L.J., Hamilton, M.A., Horan, M.J. and Scoates, R.F.J., 2004, U-Pb zircon and Re-Os isotope geochronology of mineralized ultramafic intrusions and associated nickel ores from the Thompson Nickel Belt, Manitoba, Canada: *Economic Geology* (in press).
- Irving, E., Baker, J., Hamilton, M. and Wynne, P.J., 2004, Early Paleoproterozoic geomagnetic field in western Laurentia: implications for paleolatitude, local rotations and stratigraphy: *Precambrian Research*, v. 129, p. 251-270.
- Isley, A. E. and Abbott, D. H., 2002, Implications for the temporal distribution of high-Mg magmas for mantle plume volcanism through time: *The Journal of Geology*, v. 110, p. 141-158.
- Jackson, G.D. and Taylor, F.C., 1972, Correlation of major Archean rock units in the northeastern Canadian Shield: *Canadian Journal of Earth Sciences*, v. 9, p. 1650-1669.
- James, D.T., Kamo, S. and Krogh, T., 2002, Evolution of 3.1 and 3.0 Ga volcanic belts and a new thermotectonic model for the Hopedale Block, North Atlantic craton (Canada): *Canadian Journal of Earth Sciences*, v. 39, p. 687-710.
- Johnston, S.T., Wynne, P.J., Francis, D., Hart, C.J.R., Enkin, R.J. and Engebretson, D.C., 1996, Yellowstone in Yukon: the Late Cretaceous Carmacks Group: *Geology*, v. 24, p. 997-1000.
- Ketchum, J.W.F., Jackson, S.E., Culshaw, N.G. and Barr, S.M., 2001, Depositional and tectonic setting of the Paleoproterozoic Lower Aillik Group, Makkovik, Province, Canada: evolution of a passive margin-foredeep sequence based on petrochemistry and U-Pb (TIMS and LAM-ICP-MS) geochronology: *Precambrian Research*, v. 105, p. 331-356.
- Keppie, J.D. and Dostal, J., 1994, Late Silurian – Early Devonian transpressional rift origin of the Quebec Reentrant, northern Appalachians: constraints from geochemistry of volcanic rocks: *Tectonics*, v. 13, p. 1183-1189.
- Keppie, J.D. and Krogh, T.E., 2000, 440 Ma igneous activity in the Meguma Terrane, Nova Scotia, Canada: part of the Appalachian overstep sequence: *American Journal of Science*, v. 300, p. 528-538.
- LeCheminant, A.N. and Heaman, L.M., 1989, Mackenzie igneous events, Canada: Middle Proterozoic hotspot magmatism associated with ocean opening: *Earth and Planetary Science Letters*, v. 96, p. 38-48.
- LeCheminant, A.N., Heaman, L.M., van Breemen, O., Ernst, R.E., Baragar, W.R.A. and Buchan, K.L., 1996, Mafic magmatism, mantle roots and kimberlites in the Slave craton, in LeCheminant, A.N., et. al. (eds.), *Searching for Diamonds in Canada: Geological Survey of Canada Open File 3228*, p. 161-169.
- LeCheminant, A.N., Buchan, K.L., van Breemen, O. and Heaman, L.M., 1997, Paleoproterozoic continental break-up and reassembly: evidence from 2.19 Ga diabase dyke swarms in the Slave and western Churchill provinces, Canada: [abstract] Abstract Volume 22, Geological Association of Canada/Mineralogical Association of Canada Annual Meeting, Ottawa '97, May 19-21, 1997, p. A86.
- Legault, F., Francis, D., Hynes, A. and Budkewitsch, P., 1994, Proterozoic continental volcanism in the Belcher Islands: Implications for the evolution of the Circum Ungava Fold Belt: *Canadian Journal of Earth Sciences*, v. 31, p. 1536-1549.
- MacDonald, L.A., Barr, S.M., White, C.E., and Ketchum, J.W.F., 2002, Petrology, age, and tectonic setting of the White Rock Formation, Meguma terrane, Nova Scotia: evidence for Silurian continental rifting: *Canadian Journal of Earth Sciences*, v. 39, p. 259-277.
- Maher, H.D., 2001, Manifestations of the Cretaceous High Arctic Large Igneous Province in Svalbard: *The Journal of Geology*, v. 109, p. 91-104.
- Marillier, F. and Verhoef, J., 1989, Crustal thickness under the Gulf of St. Lawrence, northern Appalachians, from gravity and deep seismic data: *Canadian Journal of Earth Sciences*, v. 26, p. 1517-1532.
- Maurice, C., Francis, D. and Madore, L., 2003, Constraints on early Archean crustal extraction and tholeiitic-komatiitic volcanism in greenstone belts of the Northern Superior Province: *Canadian Journal of Earth Sciences*, v. 40, p. 431-445.
- McHone, J.G., 1996, Constraints on the mantle plume model for Mesozoic alkaline intrusions in northeastern North America: *Canadian Mineralogist*, v. 34, p. 325-334.
- Modeland, S., Francis, D. and Hynes, A., 2003, Enriched mantle components in Proterozoic continental-flood basalts of the Cape Smith foldbelt, northern Québec: *Lithos*, v. 71, p. 1-17.

- Moore, E.M., 2002, Pre-1 Ga (pre-Rodinian) ophiolites: their tectonic and environmental implications: Geological Society of America, Bulletin, v. 114, p. 80-95.
- Murphy J.B., van Staal, C.R. and Keppie, J.D., 1999, Middle to late Paleozoic Acadian orogeny in the northern Appalachians: A Laramide-style plume-modified orogeny?: *Geology*, v. 27, p. 653-656.
- Murphy, J.B., Hynes, A.J., Johnston, S.T. and Keppie, J.D., 2003, Reconstructing the ancestral Yellowstone plume from accreted seamounts and its relationship to flat-slab subduction: *Tectonophysics*, v. 365, p. 185-194.
- Naldrett, A.J., 1999, World-class Ni-Cu-PGE deposits: key factors in their genesis: *Mineralium Deposita*, v. 34, p. 227-240.
- Ojakangas, R.W., Morey, G.B. and Green, J.C., 2001, The Mesoproterozoic Midcontinent rift system, Lake Superior region, USA: *Sedimentary Geology*, v. 141-142, p. 421-442.
- Pálffy, J., 2003, Volcanism of the Central Atlantic Magmatic Province as a potential driving force in the end-Triassic mass extinction, *in* Hames, W.E., McHone, J.G., Renne, P.R., and Ruppel, C. (eds.), *The Central Atlantic Magmatic Province: insights from fragments of Pangea*: American Geophysical Union, Geophysical Monograph, v. 136, p. 255-267.
- Park, J.K., Buchan, K.L., and Harlan, S.S., 1995, A proposed giant radiating dyke swarm fragmented by the separation of Laurentia and Australia – based on paleomagnetism of circa 780 Ma mafic intrusions in western North America: *Earth and Planetary Science Letters*, v. 132, p. 129-139.
- Pe-Piper, G. and Piper D.J.W., 1998, Geochemical evolution of Devonian-Carboniferous igneous rocks of the Magdalen basin, eastern Canada: Pb- and Nd-isotope evidence for mantle and lower crustal sources: *Canadian Journal of Earth Sciences*, v. 35, p. 201-221.
- Percival, J.A., Bleeker, W., Cook, F.A., Rivers, T., Ross, G. and van Staal, C., 2004, PanLITHOPROBE Workshop IV: Intra-orogen correlations and comparative orogenic anatomy: *Geoscience Canada*, v. 31, p. 23-39.
- Peterson, T.D., Van Breemen, O., Sandeman, H. and Cousens, B. 2002. Proterozoic (1.85-1.75 Ga) igneous suites of the Western Churchill Province: granitoid and ultrapotassic magmatism in a reworked Archean hinterland. *Precambrian Research*, v. 119, p. 73-100.
- Pirajno, F., 2000, *Ore Deposits and Mantle Plumes*: Kluwer Academic, Dordrecht, 556 p.
- Prokoph, A., Ernst, R.E., Buchan, K.L. and El Bilali, H., 2003, Linkage between emplacement of large igneous provinces and seawater composition through the past 3500 million years [abstract]: *in* Kerr, A., England, R. and Wignall, P. (convenors), *Mantle plumes: Physical processes, chemical signatures, biological effects*. Multidisciplinary meeting held at Cardiff University and the National Museum and Gallery, Cardiff, Wales, 11-12 September 2003.
- Prokoph, A., Ernst, R.E. and Buchan, K.L., 2004, Time-series analysis of Large Igneous Provinces: 3500 Ma to Present: *The Journal of Geology*, v. 112, p. 1-22.
- Puffer, J.H., 2002, A Late Neoproterozoic eastern Laurentian superplume: location, size, chemical composition, and environmental impact: *American Journal of Science*, v. 302, p. 1-27.
- Rainbird, R. and Ernst, R.E., 2001, The sedimentary record of mantle-plume uplift, *in* Ernst, R.E. and Buchan, K.L. (eds.), *Mantle Plumes: Their Identification Through Time*: Geological Society of America, Special Paper 352, Boulder, CO., p. 227-245.
- Reichow, M.K., Saunders, A.D., White, R.V., Pringle, M.S., Al-Mukhamedov, A.I., Medvedev, A.I. and Kirda, N.P., 2002, ⁴⁰Ar/³⁹Ar dates from the West Siberian Basin: Siberian flood basalt province doubled: *Science*, v. 296, p. 1846-1849.
- Richards, M.A., Jones, D.L., Duncan, R.A. and De Paolo, D.J., 1991, A mantle plume initiation model for the Wrangellia flood basalt and other oceanic plateaus: *Science*, v. 254, p. 263-267.
- Rivers, T. and Corrigan, D., 2000, Convergent margin on southeastern Laurentia during the Mesoproterozoic: tectonic implications: *Canadian Journal of Earth Sciences*, v. 37, p. 359-383.
- Rock, N.M.S., 1991, *Lamprophyres*: Blackie and Co. Publishers Ltd., Glasgow, United Kingdom, 285 p.
- Ross, G.M. and Eaton, D.W., 1997, Winagami reflection sequence: seismic evidence for postcollisional magmatism in the Proterozoic of western Canada: *Geology*, v. 25, p. 199-202.
- Ryan, B. and James, D. 2004. The Mesoproterozoic Nain Plutonic Suite and its country rocks in the Kingurutik Lake-Fraser River area, Labrador (NTS 14D/9 and 16), *in* Current Research, Newfoundland Department of Mines and Energy, Geological Survey, Report 04-1, p. 235-258.
- Sanborn-Barrie, M., and Skulski, T., 1999, Tectonic assembly of continental margin and oceanic terranes at 2.7 Ga in the Savant Lake – Sturgeon Lake greenstone belt, Ontario: *in* Current Research Part C; Geological Survey of Canada, Paper 1999-C, p. 209-220.
- Schau, M., 1997, Geology of the Archean Prince Albert Group in the Richards Bay area, northeastern Melville Peninsula, District of Franklin, Northwest Territories: Geological Survey of Canada, Bulletin 385, 44 p.
- Schneider, D.A., Bickford, M.E., Cannon, W.F., Schulz, K.J. and Hamilton, M.A., 2002, Age of volcanic rocks and syndepositional iron formations, Marquette Range Supergroup: implications for the tectonic setting of Paleoproterozoic iron formations of the Lake Superior region: *Canadian Journal of Earth Sciences*, v. 39, p. 999-1012.
- Schwarz, E.J. and Fujiwara, Y. 1981. Paleomagnetism of the circum-Ungava fold belt II: Proterozoic rocks of Richmond Gulf and Manitounuk Islands, *in* Campbell, F.H.A. (ed.) *Proterozoic Basins of Canada*: Geological Survey of Canada, Paper 81-10, p. 255-267.
- Schwab, D.L., Thorkelson, D.J., Mortensen, J.K., Creaser, R.A. and Abbott, J.G., 2004, The Bear River dykes (1265-1269 Ma) extend the Mackenzie swarm into the Yukon, Canada: *Precambrian Research* (in press).
- Schissel, D. and Smal, R., 2001, Deep-mantle plumes and ore deposits: *in* Ernst, R.E. and Buchan, K.L. (eds.), *Mantle Plumes: Their Identification Through Time*: Geological Society of America, Special Paper 352, Boulder CO., p. 291-322.
- Scott, D.J., St-Onge, M.R., Lucas, S.B. and Helmstaedt, H., 1999, The 2.00 Ga Purtunig ophiolite, Cape Smith Belt, Canada: MORB-like crust intruded by OIB-like magmatism: *Ophiolite*, v. 24, p. 199-215.
- Scott, D.J., St-Onge, M.R. and Corrigan, D., 2002, Geology of the Paleoproterozoic Piling Group and underlying Archean gneiss, central Baffin Island, Nunavut: *in* Current Research, Part C; Geological Survey of Canada, Paper 2002-C17, 10 p.
- Sears, J.W., Price, R.A. and Khudoley, A.K., 2004, Linking the Mesoproterozoic Belt-Purcell and Udzha basins across the west Laurentia – Siberia connection: *Precambrian Research*, v. 129, p. 291-308.
- Şengör, A.M.C., 2001, Elevation as indicator of mantle-plume activity, *in* Ernst, R.E. and Buchan, K.L., eds., *Mantle Plumes: Their Identification Through Time*: Geological Society of America, Special Paper 352, Boulder CO., p. 183-225.
- Sevigny, J.H., Cook, F.A. and Clark, E.A., 1991, Geochemical signature and seismic stratigraphic setting of Coppermine basalts drilled beneath the Anderson Plains in Northwest Canada: *Canadian Journal of Earth Sciences*, v. 28, p. 184-194.
- Skulski, T.M., Sandeman, H., MacHattie, T., Sanborn-Barrie, M., Rayner, N., and Byrne, D., 2003a, Tectonic setting of the 2.73-2.70 Ga Prince Albert Group, Rae domain, Nunavut: [abstract] *in* Vancouver 2003, Joint meeting of the Geological Association of Canada, Mineralogical Association of Canada and the Society of Exploration Geophysicists, 25-28 May, 2003.
- Skulski, T., Sandeman, H., Sanborn-Barrie, M., MacHattie, T., Young, M., Carson, C.,

- Berman, R., Brown, J., Rayner, N., Panagapko, D., Byrne, D. and Deyell, C., 2003b, Bedrock geology of the Ellice Hills map area and new constraints on the regional geology of the Committee Bay area, Nunavut: *in* Current Research, Part C; Geological Survey of Canada, Paper 2003-C22.
- Sproule, R.A., Leshner, C.M., Ayer, J.A., Thurston, P.C. and Herzberg, C.T., 2002, Spatial and temporal variations in the geochemistry of komatiites and komatiitic basalts in the Abitibi greenstone belt: *Precambrian Research*, v. 115, p. 153-186.
- Stern, R.A., Machado, N., Syme, E.C., Lucas, S.B. and David, J., 1999, Chronology of crustal growth and recycling in the Paleoproterozoic Amisk collage (Flin Flon Belt), Trans-Hudson orogen, Canada: *Canadian Journal of Earth Sciences*, v. 36, p. 1807-1827.
- Stott, G.M. and Halls, H.C., 2002, Paleomagnetic, geochemical and U/Pb geochronologic studies of mafic dikes in northern Ontario: Relevance to mineralization associated with the Nipigon embayment and kimberlites, in Baker, C.L., Debicki, E.J., Kelly, R.I. and Parker, J.R. eds., Summary of Field Work and Other Activities 2002, Ontario Geological Survey, Open File Report 6100, pp. 13-1 to 13-10.
- St-Onge, M.R., Scott, D.J. and Lucas, S.B., 2000, Early partitioning of Quebec: microcontinent formation in the Paleoproterozoic: *Geology*, v. 28, p. 323-326.
- Struik, L.C., Schiarizza, P., Orchard, M.J., Cordey, F., Sano, H., MacIntyre, D.G., Lapierre, H. and Tardy, M., 2001, Imbricate architecture of the upper Paleozoic to Jurassic oceanic Cache Creek Terrane, central British Columbia: *Canadian Journal of Earth Sciences*, v. 38, p. 495-514.
- Syme, E.C., Lucas, S.B., Bailes, A.H. and Stern, R.A., 1999, Contrasting arc and MORB-like assemblages in the Paleoproterozoic Flin Flon belt, Manitoba, and the role of intra-arc extension in localizing volcanic-hosted massive sulphide deposits: *Canadian Journal of Earth Sciences*, v. 36, p. 1767-1788.
- Stubley, M., 2003, Spatial distribution of kimberlite in the Slave craton: a geometrical approach: [extended abstract], 8th International Kimberlite Conference, Vancouver, Canada.
- Tarduno, J.A., Brinkman, D.B., Renne, P.R., Cottrell, R.D., Scher, H., and Castillo, P., 1998, Evidence for extreme climatic warmth from Late Cretaceous Arctic vertebrates: *Science*, v. 282, p. 2241-2243.
- Tardy, M., Lapierre, H., Struik, L.C., Bosch, D. and Brunet, P., 2001, The influence of mantle plume in the genesis of the Cache Creek oceanic igneous rocks: implications for the geodynamic evolution of the inner accreted terranes of the Canadian Cordillera: *Canadian Journal of Earth Sciences*, v. 38, p. 515-534.
- Thorkelson, D.J., Mortensen, J.K., Creaser, R.A., Davidson, G.J. and Abbott, J.G., 2001, Early Proterozoic magmatism in Yukon, Canada: constraints on the evolution of northwestern Laurentia: *Canadian Journal of Earth Sciences*, v. 38, p. 1479-1494.
- Thorkelson, D.J., Abbott, J.G., Roots, C.F. and Gordey, S.P., 2003, Early Proterozoic to early Paleozoic supracrustal history of Yukon: in Vancouver, 2003, Joint meeting of the Geological Association of Canada-Mineralogical Association of Canada-Society of Exploration Geophysicists 2003, Abstract 107 (CD-ROM and print version)
- Tomlinson, K.Y. and Condie, K.C., 2001, Archean mantle plumes: evidence from greenstone belt geochemistry, *in* Ernst, R.E. and Buchan, K.L., (eds.), Mantle Plumes: Their Identification Through Time: Geological Society of America, Special Paper 352, p. 341-357.
- Tomlinson, K.Y., Stevenson, R.K., Hughes, D.J., Hall, R.P. Thurston, P.C. and Henry, P., 1998, The Red Lake greenstone belt, Superior province: evidence of plume-related magmatism at 3 Ga and evidence of an older enriched source: *Precambrian Research*, v. 89, p. 59-76.
- Tomlinson, K.Y., Hughes, D.J., Thurston, P.C. and Hall, R.P., 1999, Plume magmatism and crustal growth at 2.9 to 3.0 Ga in the Steep Rock and Lumby Lake area, western Superior Province: *Lithos*, v. 146, p. 103-136.
- Turek A., Woodhead, J. and Zwanzig, H.V., 2000, U-Pb age of the gabbro and other plutons at Lynn Lake (part of NTS 64C): *in* Report of Activities 2000, Manitoba Industry, Trade and Mines, Manitoba Geological Survey, p. 97-104.
- Turner, W.A., Heaman, L.M. and Creaser, R.A., 2003, Sm-Nd fluorite dating of Proterozoic low-sulfidation epithermal Au-Ag deposits and U-Pb zircon dating of host rocks at Mallery Lake, Nunavut, Canada: *Canadian Journal of Earth Sciences*, v. 40, p. 1789-1804.
- Trettin, H.P. and Parrish, R., 1987, Late Cretaceous bimodal magmatism, northern Ellesmere Island: isotopic age and origin: *Canadian Journal of Earth Sciences*, v. 24, p. 257-265.
- Ueng, W.C., Fox, T.P., Larue, D.K. and Wilband, J.T., 1988, Geochemistry and petrogenesis of early Proterozoic Hemlock volcanic rocks and the Kiernan sills, southern Lake Superior region: *Canadian Journal of Earth Sciences*, v. 25, p. 528-546.
- Upton, B.G.J., Emeleus, C.H., Heaman, L.M., Goodenough, K.M., and Finch, A.A., 2003, Magmatism of the mid-Proterozoic Gardar Province, South Greenland: chronology, petrogenesis and geological setting: *Lithos*, v. 68, p. 43-65.
- van Staal, C.R., Sullivan, R.W. and Whalen, J.B., 1996, Provenance and tectonic history of the Gander Zone in the Caledonian/Appalachian orogen: implications for the origin and assembly of Avalon, *in* Nance, R.D. and Thompson, M.D. (eds.), Avalonian and Related Peri-Gondwanan Terranes of the Circum-North Atlantic: Geological Society of America Special Paper 304, Boulder, Colorado.
- Van Wagoner, N., Leybourne, M.I., Dadd, K.A. and Huskins, M.L.A., 2001, The Silurian(?) Passamaquoddy Bay mafic dyke swarm, New Brunswick: petrogenesis and tectonic implications: *Canadian Journal of Earth Sciences*, v. 38, p. 1565-1578.
- Wardle, R.J., James, D.T, Scott, D.J. and Hall, J., 2002, The southeastern Churchill Province: synthesis of a Paleoproterozoic transpressional orogen: *Canadian Journal of Earth Sciences*, v. 39, p. 639-663.
- Waldron, J.W.F. and van Staal, C.R., 2001, Taconic orogeny and the accretion of the Dashwoods block: a peri-Laurentian microcontinent in the Iapetus Ocean: *Geology* v. 29, p. 811-814.
- White, R.S. and McKenzie, D.P., 1989, Magmatism at rift zones: the generation of volcanic continental margins and flood basalts: *Journal of Geophysical Research*, v. 94, p. 7685-7729.
- Wilkinson, L., Kjarsgaard, B.A., LeCheminant, A.N. and Harris, J., 2001, Diabase dyke swarms in the Lac de Gras area, Northwest Territories and their significance to kimberlite exploration: initial results: *in* Current Research, Part C; Geological Survey of Canada, Paper 2001-C8, 17 p.
- Wyman DA., 1999, A 2.7 Ga depleted tholeiitic suite: evidence of plume-arc interaction in the Abitibi belt, Canada: *Precambrian Research*, v. 97, p. 27-42
- Zaleski, E., Davis, W.J. and Sandeman, H.A., 2001, Continental extension, mantle magmas and basement cover relationships, *in* Cassidy, K.F., Dumphy, J.M. and Van Kranendonk, M.J. eds., extended abstracts, 4th International Archean Symposium 2001, Australian Geological Survey Organization - Geoscience Australia, Record 2001/37, p.374-376.
- Zwanzig, H.V., Bailes, A.H. and Böhm, Ch.O., 2001, Josland Lake sills: U-Pb age and tectonostratigraphic implications (parts of NTS 63K and 63N): *in* Report of Activities 2001, Manitoba Industry, Trade and Mines, Manitoba Geological Survey, p. 28-32.

Geological Survey of Canada publication no. 2004026

Accepted as revised 30 June 2004