

The Tooth of Time: Charlie Roots

Paul F. Hoffman

Volume 39, numéro 4, 2012

URI : https://id.erudit.org/iderudit/geocan39_4clm01

[Aller au sommaire du numéro](#)

Éditeur(s)

The Geological Association of Canada

ISSN

0315-0941 (imprimé)

1911-4850 (numérique)

[Découvrir la revue](#)

Citer cet article

Hoffman, P. F. (2012). The Tooth of Time:: Charlie Roots. *Geoscience Canada*, 39(4), 185–194.

COLUMN

The Tooth of Time: Charlie Roots

Paul F. Hoffman

1216 Montrose Avenue,
Victoria, British Columbia
V8T 2K4

Charlie Roots is a 56 year old field geologist with the Geological Survey of Canada (GSC) who lives in Whitehorse and is based at the Yukon Geological Survey, a territorial government organization. This is the story of my three brief encounters with Charlie, each separated by an hiatus of more than 15 years. Our most important and recent encounter was wordless—no meeting, no phone call, no email—the perfect crime. Now, I must admit at the outset that when I learned in school that much of the Yukon had ‘escaped’ Pleistocene glaciation, I wrote it off for my purposes as an area of good soil and bad outcrop. I was mistaken.

In January 1977, I was in a bit of a jam. The previous summer, with a crew of six, I had mapped the East Arm of Great Slave Lake, NWT, a system of deep channels and bays extending 240 km into the Precambrian Shield (Fig. 1). Christie Bay, the third deepest column of freshwater in the world after lakes Baikal and Victoria, owes its existence to erosion by ice streams within the Laurentide Ice Sheet, guided by a canoe-shaped synclinorium of soft, nearly unmetamorphosed, sedimentary and associated igneous rocks of Orosirian (late Paleoproterozoic, 2.05-1.80 Ga) age. As mapped by Cliff Stockwell (Stockwell 1932, 1936), the synclinorium is a doubly-plunging upright structure, more tightly folded on its south-eastern side, where it is broken by




Figure 1. Oblique ESE-looking satellite view of the East Arm of Great Slave Lake, Northwest Territories. Christie Bay, the third deepest freshwater column in the world, was carved by Laurentide ice streams flowing toward the lower right. The East Arm is underlain by nearly unmetamorphosed Orosirian (2.05-1.80 Ga) sedimentary and igneous rocks, deformed by thin-skinned thrusts (not shown) and a system of right-slip faults (dashed lines) related to collisional indentation of Slave craton into Rae craton. Great Slave Lake shear zone is a dextral mylonite zone developed within a magmatic arc on the leading edge of Rae craton (Hoffman 1987). Reddish patches are recent forest fires.

faults. One of them, the McDonald Fault, presents a linear escarpment that parallels the east arm and puts granitoid basement rocks against the soft sedimentary succession. Stockwell's map accurately portrays the geology that he saw on the shorelines of the East Arm and accessible waterways to the south (Stark Lake, Murky Channel), but without aircraft support he was

unable to map all the rocks as far south as the McDonald Fault. My thesis project on the sedimentary succession revealed only one flaw in Stockwell's map: a major carbonate platform and ramp, the Pethei Group, had a demonstrably-isochronous basinal facies (Hoffman 1974) that he mistakenly assigned to the stratigraphically overlying Stark Formation. Following

the discovery of an Orosirian continental margin in the Wopmay orogen to the west, to which the main part of the sedimentary succession in the East Arm is genetically related (Hoffman 1969), I had disastrously interpreted the foldbelt as an 'aulacogen', the failed arm of a rift system active at the creation of the continental margin, analogous to the Cenozoic Ethiopian Rift Valley, or the subsided Early Cretaceous rift arm at the head of the Niger Delta (Hoffman 1973).

The urge to systematically remap the East Arm arose during a year of teaching in the University of California at Santa Barbara, on unpaid leave from GSC. There, John Crowell introduced me to the San Andreas strike-slip fault system and related sedimentary basins (Wallace 1990). He had demonstrated not only the large displacement on the fault but also its strictly Neogene age (Crowell 1962), surprising many who thought that great faults exploit ancient lines of crustal weakness. In map pattern, the McDonald Fault system (Stockwell 1936) has strong similarities and significant differences compared with the San Andreas—it is a *left-stepping* right-slip system, linked not by rhomboclasts (Carey 1958) but rhombic pop-up structures. Large strike-slip displacements (Thomas et al. 1976) would have to be estimated in order to restore the Orosirian sedimentary basin. An off season between three-year, 1:250,000-scale, mapping projects in northern Wopmay orogen offered the opportunity to remap the East Arm in 1976, my 10-year-old thesis project having been reactivated by GSC for this purpose. Straight away we found, between the East Arm and the McDonald Fault, a stack of recurrently-folded basinal-facies thrust nappes, refolded by the doubly-plunging upright folds (Hoffman et al. 1977). The refolded nappes extend for 200 km along strike. They were the first nail in the coffin of the aulacogen concept as applied to the East Arm of Great Slave Lake, which eventually emerged as a collisional indentation structure analogous to the Cenozoic Bay of Bengal – Indo-Burman Ranges area of the India-Eurasia collision zone (Gibb and Thomas 1977; Badham 1978; Gibb 1978; Hoffman




Figure 2. Mosaic of 1:50,000-scale geological maps covering the East Arm of Great Slave Lake, Northwest Territories, hand-coloured by Charlie Roots for display at the 1997 GAC-MAC annual meeting in Vancouver and published as GSC Open File 475. Photographed using period equipment by direct sunlight in Victoria, BC, in December, 2012. For legend, see GSC Map 1628A (Hoffman 1980) available as a free .pdf download from the Geoscan database: <http://geoscan.ess.nrcan.gc.ca/starweb/servlet.starweb>

1987; Johnson 1990; Ritts and Grotzinger 1994). A host of other surprises emerged in the 19 weeks (and 1500 gallons of 'kicker' gas) it took to remap the East Arm. In a lifetime of geological travels, I never again encountered such a diversity of geological phenomena, exquisitely exposed, in so small an area (Fig. 2).

The following January, a preliminary report on the East Arm was published (Hoffman et al. 1977) and compilation of twenty-five 1:50,000-scale geological maps (Hoffman 1977) was well underway. I hoped to display the maps at the GAC-MAC meeting in May, before embarking on the next mapping project in Wopmay orogen. I could finish the maps if I suspended my extracurriculars (e.g. running), but would still need help hand-colouring them for display—this was before digital map production and colour printers. The Winter Works program, a federal Liberal government initiative to reduce seasonal youth unemployment, provided help. Ira Stevenson, a congenial geologist who had mapped in Nova Scotia, northern Quebec and Labrador, managed the program at GSC. On cue, he came down to my office and introduced a broad-shouldered young man

with sandy-red hair and a snaggle-toothed grin. Charlie Roots was a third-year student from Ottawa at Dartmouth College in Hanover, NH, taking a semester break from classes. "Charlie is Fred Roots' boy," Ira told me by way of introduction, "Fred Roots of the Polar Continental Shelf Program." In the wake of *Sputnik*, PCSP was launched in 1958 as a federal government program to encourage international scientific research throughout the Canadian Arctic by providing communications, equipment, accommodations, liaison with residents and air support at contract rates. PCSP was Fred Roots' brainchild and he directed it for 14 years before becoming Science advisor to the federal government Minister of the Environment at the department's inception in 1972. He began as a geologist. When the Norwegian-British-Swedish Antarctic Expedition of 1949-52 explored Queen Maud Land, conjugate to the east coast of southern Africa, the only thing British about the chief geologist was British Columbia, where Fred Roots was born and educated (BSc, MSc – UBC; PhD – Princeton). As the first modern and overtly international scientific expedition in the polar

regions without territorial pretensions, it prepared the way for the Third International Polar Year, which became the International Geophysical Year, 1957-58 (Korsmo 2007; Roots 2011). In 1964, Fred Roots led the first Canadian Himalayan Expedition, which named and climbed high on an elegant pyramidal peak framed by glaciers, Sang-E-Marmar, which is nestled in the southern Batura Range (Figs. 3, 4) facing the Hunza Valley, in the Karakoram Mountains of northern Pakistan (Searle 1991). Whether as Founding Chairman of the International Arctic Science Committee or Canadian representative to the Antarctic Treaty, Fred Roots was a humble and soft-spoken bulldog for science and international science in the Polar regions.

Yet, Charlie Roots' passion for high places was inspired by Noye Johnson at Dartmouth College. Johnson, who died in 1987 at the age of 57, was a remarkably versatile, field-oriented hydrogeologist, low-temperature geochemist, geochronologist, Cenozoic terrestrial stratigrapher and mammalian paleontologist (Naeser and Zeitler 1990). Starting in 1963, his systematic long-term study of a forested watershed-ecosystem in the White Mountains of New Hampshire yielded seminal insights on geochemical cycling, weathering rates and, coincidentally, 'acid rain' (Likens, Bormann and Johnson 1972). His interest in geochronology began even earlier with basic research on thermoluminescence, followed by regional-scale magnetostratigraphy and fission-track dating of detrital grains in Cenozoic sedimentary basins in active tectonic areas of Pakistan, Bolivia, Argentina and the southwestern United States. Above all, Noye Johnson was a man of action. Charlie was taking the semester off to work along side an Inuit contemporary, Zacharias Kunnuk from Igloolik, as labourers. Their job was to dig out an ice-core drilling camp (buried by snow) on the Devon Island Ice Cap in April, move it 350 km to the north in eight Twin Otter loads, rebuild it and core the Agassiz Ice Cap on Ellesmere Island down to bedrock. Later the same year, he would participate in reconnaissance mapping of Canada's 'Himalaya', the St. Elias Mountains of




Figure 3. Sang-E-Marmar (6,949 m), center, viewed from the south, was named by the First Canadian Himalayan Expedition in 1964 led by E.F. Roots. The outlet valley (foreground) drains into the middle Hunza River valley in the Karakoram Mountains northeast of Gilgit. The expedition intended to locate and climb Hachindar Chhish (upper left), which proved to be too technically difficult to attempt. They reached 6,300 m on Sang-E-Marmar before being forced to retreat by heavy snowstorms. The peak was first climbed in 1984 by a team from Osaka University, Japan. According to Searle (1991), a southwest-directed thrust on the lower flank of Sang-E-Marmar places Cretaceous (95 Ma) Hunza granodiorite over Paleozoic-Early Mesozoic marble (Dumordu unit), for which the mountain is named. Pasu Sar (7,478 m) at top was first climbed in 1994.



Figure 4. Dissected alluvial fan, near Askole on the Braldu River ~110 km southeast of Sang-E-Marmar, is a testament to rapid exhumation and uplift of the Karakoram Mountains. Note the agricultural terraces, irrigated by an aqueduct (upper left) tapping glacial meltwater, on which the region depends. Photo in 1985 by the author.

SW Yukon (Fig. 5), when Kluane National Park was created. If he felt over-qualified for his Winter Works job with me—colouring a 5 x 3 metre map (Fig. 2) by hand with colour pencils—he didn't say so.

The Spring on the eastern Arctic ice caps and Summer in the St. Elias Mountains set Charlie's compass permanently on North. Returning to Ottawa for his PhD at Carleton University with John Moore and Al Donaldson, Charlie studied the Neoproterozoic Mount Harper Volcanic Complex as part of GSC's Ogilvie Mountains Project (Thompson and Roots 1982). As a rift-related bimodal shield volcano associated with the breakup of Rodinia, the Mount Harper Group is unique on the Cordilleran margin of Laurentia (Roots 1987; Mustard and Roots 1997). His thesis won the Léopold Gélinas Award of the Volcanology Division of GAC. In 1987, Charlie moved to the Cordilleran Division of GSC in Vancouver, where he worked with Bob Thompson and Peter Mustard mapping the Dawson 1:250K sheet in west-central Yukon (Thompson et al. 1994). They enlisted geochronologist Randy Parrish in Ottawa, who obtained a rather imprecise U-Pb zircon date of $751 \pm 26/-18$ Ma for a rhyolite lava near the top of the volcanic complex (Roots and Parrish 1988). Working with paleomagnetist John Park, Charlie puzzled over evidence that the volcanic complex was rotated $\sim 90^\circ$ anticlockwise around a near-vertical axis, compared with statistically coeval 780-Ma intrusions in the Mackenzie Mountains and the Canadian Shield (Park et al. 1992, 1995). He encouraged Peter Mustard to study the sedimentary rift-basin that preceded Mount Harper volcanism (Mustard 1991; Mustard and Roots 1997). Charlie pecked away at 1:50,000-scale mapping in the Ogilvie Mountains whenever funds were available.

In 1992, Charlie joined a 16-person expedition to climb and place a GPS receiver at the top of Mount Logan, to accurately determine the height of Canada's tallest mountain as part of the 150th anniversary celebrations of GSC (Schmidt 1993). I had just left the organization and wasn't in a celebratory mood when I met Charlie



Figure 5. Kaskawulsh Glacier, looking southwest into the St. Elias Mountains, south of Kluane Lake. This was part of the area mapped in reconnaissance by Dick Campbell (GSC) assisted by Charlie Roots in 1977 when Kluane National Park was established. Note the complex of glacial and proglacial deposits in the center foreground.

at the Cordilleran Roundup in Vancouver in January 1993. With characteristic lack of finesse, I suggested that GSC might better be seen as having less frivolous and more geological aims than the height of a mountain. Charlie responded that under the circumstances any positive publicity was good for GSC. The GPS site was anchored in rock, so that the change in elevation of the peak over time could be measured. Ten hand samples of granite had been taken at precise points between 3400 and 5900 m elevation for apatite fission-track analysis, to estimate the cooling (exhumation) rate in an area of anomalous tectonically-driven uplift. He was asked to participate because of his familiarity with the St. Elias Mountains. Moreover, the public was more impressed by the size or shape of a mountain than by its mode of origin, whatever the scientists thought. This last point was not lost on William Logan himself. In the competition to attract immigrants waged at the 1851 World's Fair at the Crystal Palace,

London, Logan's geological display won hands down because the 'Canadian' rock specimens were neither rare nor beautiful, they were simply larger, with a strong emphasis on ores (Zeller 1987).

On October 4, 2000, Canadian Prime Minister Jean Chrétien announced that Mount Logan would be renamed Mount Pierre Elliott Trudeau after his recently-deceased predecessor. The announcement, symptomatic of the federal Liberals' inattention to western Canadian public opinion, was openly opposed by First Nation leaders, the Yukon Minister of Tourism and member of the Geographical Names Board of Canada, surviving relatives of Sir William, prominent academics, and the presidents of national geographical and geological societies. As a compromise, it was proposed that a secondary peak on the high plateau of Mount Logan be renamed instead, and Charlie Roots was consulted on this by a *National Post* reporter because of his first-hand knowledge of the mountain.

Had I known, I would have suggested they rename Mount Harper.

Charlie wrote to me a few days after our confrontation at the 1993 Cordilleran Roundup, restating his arguments. At the end of the letter, he added that *“the right conditions to propose a thesis project on the [Neoproterozoic] sedimentary succession that overlies the Wernecke Supergroup are coming around again. The succession is called the Pinguicula (Eisbacher, GSC Paper 80-27), or Fifteenmile Group (Mustard, Roots, Bob Lane work), but has never been understood nor systematically described. What is needed is a strong student, willing to unravel the structure and measure sections in a field area spanning about 200 km, and a supervisor keen on mid-Proterozoic stratigraphy and regional reconstructions. Give me a buzz if such a project tickles your fancy.”* All my resources were going to a nascent field project on the Neoproterozoic of northern Namibia and the following year I moved back east to Harvard. Years later, Noel James’ poster at the 2000 GSA meeting provided the impetus to compare Cryogenian glacial histories in Namibia and the Mackenzie Mountains, NWT (Yeo 1981; Eisbacher 1985; Aitken 1991; James et al. 2001). With U.S. federal government funding (NASA Astrobiology and NSF), we made an 18-camp reconnaissance of Cryogenian—early Ediacaran sequences and chemostratigraphy along the 500-km-long outcrop belt in the Mackenzie Mountains in three field seasons (Halverson et al. 2005, 2011; Shen et al. 2008; Hoffman and Halverson 2011). In 2004, I was accompanied by two Harvard PhDs, both of whom grew up in the Rocky Mountains. Galen Halverson (Montana ’96) was finishing his PhD on Neoproterozoic sequence and chemostratigraphy in Svalbard, Norway and Namibia; Francis Macdonald (Caltech ’01) was starting his on the Neoproterozoic-Cambrian succession in the Brooks Range of Arctic Alaska. At one camp, we densely sampled the type section of the pre-Cryogenian Little Dal Group (Gabrielse et al. 1973; Aitken 1981), a 2-km-thick carbonate-dominated sequence correlative with part of the Yukon succession Charlie had described in his letter. Finding the 0.80-Ga Bitter Springs isotope stage within the Upper Carbonate Formation




Figure 6. Distribution of Proterozoic strata in the Ogilvie, Wernecke and Mackenzie mountains of the Yukon and Northwest Territories (Macdonald et al. 2012).

of the Little Dal Group (Halverson 2006, Halverson et al. 2007) provided a target for regional correlation.

Francis Macdonald, a protégé of paleomagnetist Joe Kirschvink, had years of field experience before he started his PhD at Harvard. By the time he contacted Charlie Roots in February 2008, he had reinterpreted major Neoproterozoic successions in Arctic Alaska (Macdonald et al. 2009b), western Mongolia (Macdonald et al. 2009a) and the Tatonduk inlier (Tindir Group) astride the Yukon-Alaska international boundary (Macdonald et al. 2010a), based on aggressive field mapping and carbon isotope stratigraphy. He now proposed a regional study of Neoproterozoic (post-Wernecke Supergroup and Wernecke Breccia) and early Paleozoic rocks across the central Yukon (Fig. 6). In addition to his experience in correlative sequences to the east (Mackenzie Mountains) and west (Tatonduk inlier), he offered \$30K in NSF funding (the lion’s share of the estimated annual cost) and research collaborations. Charlie, now living in Whitehorse, responded enthusiastically. With support from Don Murphy and Maurice Colpron, he convinced Grant Abbott to approve \$5K of YGS funds for helicopter time, and GSC approved Charlie’s active involvement. Progress

reports would be published in YGS’s annual Yukon Exploration and Geology series. Field geologists being more territorial than tomcats, not everyone was happy about the new neighbours. The truth was, the hunt for a Canadian Olympic Dam deposit had showered attention on the Wernecke Supergroup and Wernecke Breccias, while few people besides Bob Thompson and Charlie had shown interest in the Neoproterozoic of the Yukon in the 15 years since his first letter.

Macdonald’s project produced immediate results. The unconformity-bounded Pinguicula, Fifteenmile and Mount Harper groups make up a carbonate-dominated early Neoproterozoic succession about 5.3 km thick, which can be correlated in detail by means of sequence and chemostratigraphy across the Yukon and with coeval successions in eastern Alaska, NWT and western Nunavut (Macdonald and Roots 2010; Macdonald et al. 2011; Halverson et al. 2012). Collaborations with biogeochemist David Johnston at Harvard and geochronologists Mark Schmitz (a Sam Bowring protégé) at Boise State University in Idaho and Alan Rooney at Durham University in the UK, produced critical dates and geochemical

tests. A U-Pb zircon date of 811.51 ± 0.25 Ma from a tuff in the upper Fifteenmile Group (Macdonald et al. 2010b) confirmed the early Neoproterozoic age of the world's oldest skeletal microfossils, the exquisitely preserved and remarkably disparate assemblage of phosphatic scales (Fig. 7) from the Tatonduk inlier (Cohen et al. 2011; Cohen and Knoll 2012). The rhyolite in the upper Mount Harper Volcanic Complex, previously dated at $751 \pm 26/-18$ Ma (Roots and Parrish 1988), was re-dated at the same site at 717.43 ± 0.14 Ma (Macdonald et al. 2010b), a striking example of ongoing improvements in the accuracy and precision of U-Pb (IDTIMS) dating. This was more than a technical display. The volcanic complex is depositionally overlapped by massive and stratified diamictite, within which a brecciated tuff in the Mount Harper area was dated at 716.47 ± 0.24 Ma (Macdonald et al. 2010b). Lonestones occur below and above the tuff horizon, and the equivalent interval in the Hart River inlier to the east of the volcanic complex contains massive diamictite bearing multiply-striated boulders and stratified diamictite with glacial plow structures, confirming Charlie's suspicion that the waning of Mount Harper volcanism coincided with the onset of Cryogenian glaciation (C.F. Roots, YGS project proposal, March 2008), equivalent to the lower Rapitan Group in the Mackenzie Mountains.

Farther afield, the dates showed that Mount Harper volcanism was coeval with the Franklin Large Igneous Province (LIP), a suite of mafic dyke swarms, sills and lavas recognized across the Canadian Arctic as far as NW Greenland (Fig. 8), dated on Victoria Island at 716.33 ± 0.54 Ma (Macdonald et al. 2010b). The highly-reliable grand-mean paleomagnetic pole for the Franklin LIP (Denyszyn et al. 2009) places the glacial and glacial marine deposits of the Rapitan Group in the Mackenzie Mountains at a paleolatitude of $18^\circ \pm 3^\circ$ (Evans and Raub 2011). By way of paleoclimatic comparison, during the Last Glacial Maximum in the Ethiopian Highlands, comparable in latitude and geographic setting, the lowest moraine formed at 3200 m above sea-level (Umer et al. 2004). The low paleolatitude for the




Figure 7. Scanning electron microscopy (SEM) images of the phosphatic scale microfossil genus *Thorakidictyon myriocanthum* (Allison and Hilgert 1986) from the upper Fifteenmile Group (mid-Neoproterozoic) of the Tatonduk inlier, Yukon Territory (from Cohen and Knoll 2012). Scale bar is 14 μ m in 1, 2 μ m in 2-3 and 5 μ m in 4. Six new genera containing 17 new species are described from macerated limestone that is 230-260 Myr older than the oldest calcareous skeletal fossils, the late Ediacaran Cloudina-Namacalathus assemblage (Macdonald et al. 2010a; Cohen et al. 2011).




Figure 8. Distribution of mafic dyke swarms, sill complexes and remnant flood basalt (Natkusiak Fm) of the ca. 717 Ma Franklin Large Igneous Province (LIP), relative to the coeval Mount Harper Volcanic Complex, Rapitan Group glacial and glacial marine deposits, and the paleoequator, based on the grand-mean paleomagnetic pole for the Franklin LIP (Denyszyn et al. 2009).




Figure 9. Global paleogeography ca. 715 Ma (modified after Hoffman and Li 2009), showing the distribution of known older Cryogenian (e.g. Rapitan Group) glacial and glacial marine formations (magenta stars) and extent of the Franklin LIP (dark brown ellipse). Note absence of high-latitude continents and location of Franklin LIP on the windward equatorial margin of Laurentia.

Rapitan glaciation means that 716.5 Ma is not just the age of a local or even regional glaciation; it is a minimum date for the onset of a *global* glaciation, either a snowball Earth (entire ocean ice covered) or a 'Jormungand' climate state, in which a sinuous band of open water straddles the equator (Abbott et al. 2011). The Franklin LIP itself was emplaced squarely across the paleoequator on the windward margin of Laurentia (Fig. 9). In a world that was already cold because the unusual preponderance of low-latitude continents maximized both silicate weathering and the planetary albedo, the eruption of large basalt plateau in an area perpetually soaked by onshore tradewinds is a plausible trigger mechanism for runaway glaciation from a climate model (GCM) perspective (Goddéris et al. 2003). This scenario is being tested by parallel Sr and Os isotope stratigraphy (Rooney et al. 2012): enhanced basalt weathering drives both systems to less radiogenic values, Os more rapidly because of the 10-100-fold difference in the residence times of Sr and Os in seawater.

Closer to home, the counterclockwise rotation of the Mount Harper paleomagnetic pole is somewhat smaller (~67°) when compared with the Franklin pole than with the 780 Ma pole (Park et al. 1992), but the tight synchronicity between the former rules out polar wander, true or otherwise. The Mount Harper Volcanic

Complex and potentially the entire Yukon Stable Block evidently underwent a large counterclockwise rotation relative to cratonic Laurentia, a rotation that has long been conjectured (Eisbacher 1981; Aitken and McMechan 1991; Abbott 1996) and which must be accounted for in any reconstruction of older structures, like the NNW-side-down normal faults that control early Neoproterozoic depocenters and facies in the Ogilvie Mountains (Macdonald et al. 2012). For over 40 years, Neoproterozoic rocks in the Cordillera have been viewed as the initial deposits of an early Paleozoic Cordilleran continental margin (e.g. Stewart 1972; Ross 1991), despite evidence that the margin did not exist before latest Ediacaran time (Bond and Kominz 1984). These rocks have a long and complex history, which influenced subsequent developments but cannot be explained by them (Macdonald et al. 2012). They have their own story.

YGS appears to be happy with the Neoproterozoic initiative—their contribution to Macdonald and his Harvard team's helicopter costs doubled to \$10K/year in 2009, and when Galen Halverson moved to McGill University from Adelaide in 2010, YGS provided an additional \$10K/year enabling the two teams to join forces, funded primarily by NSF (Macdonald) and NSERC (Halverson). Their field work has not been limited to the

Yukon: the Mackenzie Mountains have provided their share of excitement (Halverson et al. 2011; Johnston et al. 2012; Rooney et al. 2012). On either side of the continental divide, teams of grateful geologists camp out at Charlie and his wife Mary Ann's house in Whitehorse on their way to and from the mountains. Through relentless pursuit of research partnerships, regardless of nationality, he brought sunlight to the Neoproterozoic of central Yukon, which now glitters on the international stage.

REFERENCES

- Abbott, D.S., Voigt, A., and Koll, D., 2011, The Jormungand global climate state and implications for Neoproterozoic glaciations: *Journal of Geophysical Research*, v. 116, D18103, <http://dx.doi.org/10.1029/2011JD015927>
- Abbott, G., 1996, Implications of probable late Proterozoic dextral strike-slip movement on the Snake River fault, in *Proceedings Slave-Northern Cordillera Lithosphere Experiment, Lithoprobe Report 50*, Calgary.
- Aitken, J.D., 1981, Stratigraphy and sedimentology of the upper Proterozoic Little Dal Group, Mackenzie Mountains, Northwest Territories, in Campbell, F.H.A., ed., *Proterozoic Basins of Canada: Geological Survey of Canada, Paper 81-10*, p. 47-71.
- Aitken, J.D., 1991, The Ice Brook Formation and post-Rapitan, late Proterozoic glaciation, Mackenzie Mountains, Northwest Territories: *Geological Survey of Canada, Bulletin* 404, 43 p.
- Aitken, J.D., and McMechan, M.E., 1991, Middle Proterozoic assemblages, in Gabrielse, H., and Yorath, C., eds., *Geology of the Cordilleran Orogen in Canada*, *Geology of Canada*, v. 4, p. 97-124.
- Allison, C.W., and Hilgert, J.W., 1986, Scale microfossils from the early Cambrian of Northwest Canada: *Journal of Paleontology*, v. 60, p. 973-1015.
- Badham, J.P.N., 1978, The early history and tectonic significance of the East Arm graben, Great Slave Lake, Canada: *Tectonophysics*, v. 45, p. 201-215, [http://dx.doi.org/10.1016/0040-1951\(78\)90007-0](http://dx.doi.org/10.1016/0040-1951(78)90007-0)
- Bond, G.C., and Kominz, M.A., 1984, Construction of tectonic subsidence curves for the early Paleozoic miogeoclinal, southern Canadian Rocky Mountains: Implications for subsidence mechanisms, age of breakup,

- and crustal thinning: Geological Society of America Bulletin, v. 95, p. 155-173, [http://dx.doi.org/10.1130/0016-7606\(1984\)95<155:COTSCF>2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1984)95<155:COTSCF>2.0.CO;2)
- Carey, S.W., 1958, The tectonic approach to continental drift, in Carey, S.W., *ed.*, Continental Drift, A Symposium: Geology Department, University of Tasmania, Hobart, p. 177-355.
- Cohen, P.A., and Knoll, A.H., 2012, Scale microfossils from the mid-Neoproterozoic Fifteenmile Group, Yukon Territory: Journal of Paleontology, v. 86, p. 775-800, <http://dx.doi.org/10.1666/11-138.1>
- Cohen, P.A., Schopf, J.W., Butterfield, N.J., Kudryavtsev, A.B., and Macdonald, F.A., 2011, Phosphate biomineralization in mid-Neoproterozoic protists: Geology, v. 39, p. 539-542, <http://dx.doi.org/10.1130/G31833.1>
- Crowell, J.C., 1962, Displacement along the San Andreas fault, California: Geological Society of America, Special Paper 71, 61 p.
- Denyszyn, S.W., Halls, H.C., Davis, D.W., and Evans, D.A.D., 2009, Paleomagnetism and U-Pb geochronology of Franklin dykes in High Arctic Canada and Greenland: a revised age and paleomagnetic pole constraining block rotations in the Nares Strait region: Canadian Journal of Earth Sciences, v. 46, p. 689-705, <http://dx.doi.org/10.1139/E09-042>
- Eisbacher, G.H., 1981, Sedimentary tectonics and glacial record in the Windermere Supergroup, Mackenzie Mountains, northwestern Canada: Geological Survey of Canada, Paper 80-27, 40 p.
- Eisbacher, G.H., 1985, Late Proterozoic rifting, glacial sedimentation, and sedimentary cycles in the light of windermere deposition, western Canada: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 51, p. 231-254, [http://dx.doi.org/10.1016/0031-0182\(85\)90087-2](http://dx.doi.org/10.1016/0031-0182(85)90087-2)
- Evans, D.A.D., and Raub, T.D., 2011, Neoproterozoic glacial palaeolatitudes: a global update, in Arnaud, E., Halverson G.P., and Shields-Zhou, G., *eds.*, The Geological Record of Neoproterozoic Glaciations: Geological Society, London, Memoir 36, p. 93-112, PMID:21414169
- Gabriel, H., Blusson, S., and Roddick, J.A., 1973, Geology of the Flat River, Glacier Lake, and Wrigley Lake map-areas, District of Mackenzie and Yukon Territory: Geological Survey of Canada, Memoir 366, 421 p.
- Gibb, R.A., 1978, Slave – Churchill collision tectonics: Nature, v. 271, p. 50-52, <http://dx.doi.org/10.1038/271050a0>
- Gibb, R.A., and Thomas, M.D., 1977, The Thelon Front: A cryptic suture in the Canadian Shield?: Tectonophysics, v. 38, p. 211-222, [http://dx.doi.org/10.1016/0040-1951\(77\)90211-6](http://dx.doi.org/10.1016/0040-1951(77)90211-6)
- Goddéris, Y., Donnadieu, Y., Nédélec, A., Dupré, B., Dessert, C., Grard, A., Ramstein, G., and François, L.M., 2003, The Sturtian ‘snowball’ glaciation: fire and ice: Earth and Planetary Science Letters, v. 211, p. 1-12, [http://dx.doi.org/10.1016/S0012-821X\(03\)00197-3](http://dx.doi.org/10.1016/S0012-821X(03)00197-3)
- Halverson, G.P., 2006, A Neoproterozoic chronology, in Xiao Shuhai, and Kaufman, A.J., *eds.*, Neoproterozoic Geobiology and Paleobiology: Springer, Dordrecht, p. 231-271.
- Halverson, G.P., Hoffman, P.F., Schrag, D.P., Maloof, A.C., and Rice, A.H.N., 2005, Toward a Neoproterozoic composite carbon-isotope record: Geological Society of America Bulletin, v. 117, p. 1181-1207, <http://dx.doi.org/10.1130/B25630.1>
- Halverson, G.P., Dudás, F.Ö., Maloof, A.C., and Bowring, S.A., 2007, Evolution of the $^{87}\text{Sr}/^{86}\text{Sr}$ composition of Neoproterozoic seawater: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 256, p. 103-129, <http://dx.doi.org/10.1016/j.palaeo.2007.02.028>
- Halverson, G.P., Poitrasson, F., Hoffman, P.F., Nédélec, A., Montel, J.-M., and Kirby, J., 2011, Fe isotope and trace element geochemistry of the Neoproterozoic syn-glacial Rapitan iron formation: Earth and Planetary Science Letters, v. 309, p. 100-112, <http://dx.doi.org/10.1016/j.epsl.2011.06.021>
- Halverson, G.P., Macdonald, F.A., Strauss, J.V., Smith, E.F., Cox, G.M., and Hubert-Théou, L., 2012, Updated definition and correlation of the lower Fifteenmile Group in the central and eastern Ogilvie Mountains, in MacFarlane, K.E., and Sack, P.J., *eds.*, Yukon Exploration and Geology 2011: Yukon Geological Survey, Whitehorse, YT, p. 75-90.
- Hoffman, P.F., 1969, Proterozoic paleocurrents and depositional history of the East Arm fold belt, Great Slave Lake, Northwest Territories: Canadian Journal of Earth Sciences, v. 6, p. 441-462, <http://dx.doi.org/10.1139/e69-042>
- Hoffman, P.F., 1973, Evolution of an early Proterozoic continental margin: the Coronation geosyncline and associated aulacogens of the northwestern Canadian shield: Philosophical Transactions of the Royal Society, London, Part A, v. 273, p. 547-581.
- Hoffman, P.F., 1974, Shallow and deepwater stromatolites in an Early Proterozoic platform-to-basin facies change, Great Slave Lake, Northwest Territories: American Association of Petroleum Geologists Bulletin, v. 58, p. 856-867.
- Hoffman, P.F., 1977, Preliminary 1:50,000 scale geological maps of Athapuscow Aulacogen, East Arm of Great Slave Lake, District of Mackenzie: Geological Survey of Canada, Open File 475 (20 maps).
- Hoffman, P.F., 1980, Geology and tectonics, East Arm of Great Slave Lake, District of Mackenzie: Geological Survey of Canada, Map 1628A, 1:250,000 (geology) and 1:500,000 (tectonics).
- Hoffman, P.F., 1987, Continental transform tectonics, Great Slave Lake shear zone (ca. 1.9 Ga), northwest Canada: Geology, v. 15, p. 785-788, [http://dx.doi.org/10.1130/0091-7613\(1987\)15<785:CTTGSL>2.0.CO;2](http://dx.doi.org/10.1130/0091-7613(1987)15<785:CTTGSL>2.0.CO;2)
- Hoffman, P.F., and Halverson, G.P., 2011, Neoproterozoic glacial record in the Mackenzie Mountains, northern Canadian Cordillera, in Arnaud, E., Halverson G.P., and Shields-Zhou, G., *eds.*, The Geological Record of Neoproterozoic Glaciations: Geological Society (London), Memoir 36, p. 397-411.
- Hoffman, P.F., and Li, Zheng-Xiang, 2009, A palaeogeographic context for Neoproterozoic glaciation: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 277, p. 158-172, <http://dx.doi.org/10.1016/j.palaeo.2009.03.013>
- Hoffman, P.F., Bell, I.R., Hildebrand, R.S., and Thorstad, L., 1977, Geology of the Athapuscow Aulacogen, East Arm of Great Slave Lake, District of Mackenzie, in Report of Activities, Part A, Geological Survey of Canada, Paper 77-1A, p. 117-129.
- James, N.P., Narbonne, G.M., and Kyser, T.K., 2001, Late Neoproterozoic cap carbonates: Mackenzie Mountains, northwestern Canada: precipitation and global glacial meltdown: Canadian Journal of Earth Sciences, v. 38, p. 1229-1262, <http://dx.doi.org/10.1139/e01-046>
- Johnson, B.J., 1990, Stratigraphy and structure of the Early Proterozoic Wilson Island Group, East Arm thrust-fold belt, N.W.T.: Canadian Journal of

- Earth Sciences, v. 27, p. 552-569, <http://dx.doi.org/10.1139/e90-052>
- Johnston, D.T., Macdonald, F.A., Gill, B.C., Hoffman, P.F., and Schrag, D.P., 2012, Uncovering the Neoproterozoic carbon cycle: *Nature*, v. 483, p. 320-324, <http://dx.doi.org/10.1038/nature10854>
- Korsmo, F.L., 2007, The birth of the International Geophysical Year: *Physics Today*, v. 60, p. 38-43, <http://dx.doi.org/10.1063/1.2761801>
- Likens, G.E., Bormann, F.H., and Johnson, N.M., 1972, Acid rain: Environment: Science and Policy for Sustainable Development, v. 14, p. 33-40, <http://dx.doi.org/10.1080/00139157.1972.9933001>
- Macdonald, F.A., and Roots, C.F., 2010, Upper Fifteenmile Group in the Ogilvie Mountains and correlations of early Neoproterozoic strata in the northern Cordillera, *in* MacFarlane, K.E., Weston, L.H., and Blackburn, L.R., eds., *Yukon Exploration and Geology 2009: Yukon Geological Survey, Whitehorse, YT*, p. 237-252.
- Macdonald, F.A., Jones, D.S., and Schrag, D.P., 2009a, Stratigraphic and tectonic implications of a newly discovered glacial diamictite-cap carbonate couplet in southwestern Mongolia: *Geology*, v. 37, p. 123-126, <http://dx.doi.org/10.1130/G24797A.1>
- Macdonald, F.A., McClelland, W.C., Schrag, D.P., and Macdonald, W.P., 2009b, Neoproterozoic glaciation on a carbonate platform margin in Arctic Alaska and the origin of the North Slope subterranean: *Geological Society of America Bulletin*, v. 121, p. 448-473, <http://dx.doi.org/10.1130/B26401.1>
- Macdonald, F.A., Cohen, P.A., Dudás, F.Ö., and Schrag, D.P., 2010a, Early Neoproterozoic scale microfossils in the Lower Tindir Group of Alaska and the Yukon Territory: *Geology*, v. 38, p. 143-146, <http://dx.doi.org/10.1130/G25637.1>
- Macdonald, F.A., Schmitz, M.D., Crowley, J.L., Roots, C.F., Jones, D.S., Maloof, A.C., Strauss, J.V., Cohen, P.A., Johnston, D.T., and Schrag, D.P., 2010b, Calibrating the Cryogenian: *Science*, v. 327, p. 1241-1243, <http://dx.doi.org/10.1126/science.1183325>
- Macdonald, F.A., Smith, E.F., Strauss, J.V., Cox, G.M., Halverson, G.P., and Roots, C.F., 2011, Neoproterozoic and early Paleozoic correlations in the western Ogilvie Mountains, Yukon, *in* MacFarlane, K.E., Weston, L.H., and Relf, C., eds., *Yukon Exploration and Geology 2010: Yukon Geological Survey, Whitehorse, YT*, p. 161-182.
- Macdonald, F.A., Halverson, G.P., Strauss, J.V., Smith, E.F., Cox, G., Sperling, E.A., and Roots, C.F., 2012, Early Neoproterozoic basin formations in Yukon, Canada: Implications for the make-up and break-up of Rodinia: *Geoscience Canada*, v. 39, p. 77-99.
- Mustard, P.S., 1991, Normal faulting and alluvial-fan deposition, basal Windermere Tectonic Assemblage, Yukon, Canada: *Geological Society of America Bulletin*, v. 103, p. 1346-1364, [http://dx.doi.org/10.1130/0016-7606\(1991\)103<1346:NFAAFD>2.3.CO;2](http://dx.doi.org/10.1130/0016-7606(1991)103<1346:NFAAFD>2.3.CO;2)
- Mustard, P.S., and Roots, C.F., 1997, Rift-related volcanism, sedimentation, and tectonic setting of the Mount Harper Group, Ogilvie Mountains, Yukon Territory: *Geological Survey of Canada, Bulletin 492*, 92 p.
- Naeser, C.W., and Zeitler, P.K., 1990, In memoriam: Noye M. Johnson (1930-1987): *Journal of Geology*, v. 98, p. 423-428, <http://dx.doi.org/10.1086/629416>
- Park, J.K., Roots, C.F., and Brunet, N., 1992, Paleomagnetic evidence for rotation in the Neoproterozoic Mount Harper volcanic complex, Ogilvie Mountains, Yukon Territory, *in* Current Research, Part E: Geological Survey of Canada, Paper 92-1E, p. 1-10.
- Park, J.K., Buchan, K.L., and Harlan, S.S., 1995, A proposed giant radiating dyke swarm fragmented by the separation of Laurentia and Australia based on paleomagnetism of ca. 780 Ma mafic intrusions in western North America: *Earth and Planetary Science Letters*, v. 132, p. 129-139, [http://dx.doi.org/10.1016/0012-821X\(95\)00059-L](http://dx.doi.org/10.1016/0012-821X(95)00059-L)
- Ritts, B.D., and Grotzinger, J.P., 1994, Depositional facies and detrital composition of the Proterozoic Et-Then Group, N.W.T., Canada: sedimentary response to intracratonic indentation: *Canadian Journal of Earth Sciences*, v. 31, p. 1763-1778, <http://dx.doi.org/10.1139/e94-157>
- Rooney, A.D., Macdonald, F.A., Strauss, J.V., Dudás, F.Ö., Selby, D., Hallmann, C., 2012, Neoproterozoic glaciations and post-glacial weathering regimes: Insights from Re-Os geochronology and Os isotope stratigraphy, *in* The Neoproterozoic Era: Evolution, Glaciation, Oxygenation, 2012 Fermor Meeting, Abstract Book, Geological Society, London, p. 67.
- Roots, C.F., 1987, Regional tectonic setting and evolution of the Late Proterozoic Mount Harper Volcanic Complex, Ogilvie Mountains, Yukon: PhD thesis, Carleton University, Ottawa, ON, 180 p.
- Roots, C.F., and Parrish, R.R., 1988, Age of the Mount Harper volcanic complex, southern Ogilvie Mountains, Yukon, *in* Radiogenic Age and Isotopic Studies, Report 2: Geological Survey of Canada, Paper 88-2, p. 29-35.
- Roots, E.F., 2011, Background and evolution of some ideas and values that have led to the Antarctic Treaty, *in* Berkman, P.A., Lang, M.A., Walton, D.W.H., and Young, O.R., eds., *Science Diplomacy: Antarctica, Science, and the Governance of International Spaces*: Smithsonian Institution Scholarly Press, Washington, DC, p. 69-72, <http://dx.doi.org/10.5479/si.9781935623069.69>
- Ross, G.M., 1991, Tectonic setting of the Windermere Supergroup revisited: *Geology*, v. 19, p. 1125-1128, [http://dx.doi.org/10.1130/0091-7613\(1991\)019<1125:TSOTWS>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1991)019<1125:TSOTWS>2.3.CO;2)
- Schmidt, M., 1993, Taking GPS to the top of Canada: *GPS World*, February 1993, p. 25-34.
- Searle, M.P., 1991, *Geology and Tectonics of the Karakoram Mountains*: John Wiley & Sons, Chichester, UK, 358 p.
- Shen Yanan, Zhang Tonggang, and Hoffman, P.F., 2008, On the co-evolution of Ediacaran oceans and animals: *Proceedings of the National Academy of Sciences of the United States of America*, v. 105, p. 7376-7381, <http://dx.doi.org/10.1073/pnas.0802168105>
- Stewart, J.H., 1972, Initial deposits in the Cordilleran Geosyncline: Evidence of a Late Precambrian (<850 m.y.) continental separation: *Geological Society of America Bulletin*, v. 83, p. 1345-1360, [http://dx.doi.org/10.1130/0016-7606\(1972\)83\[1345:IDITCG\]2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1972)83[1345:IDITCG]2.0.CO;2)
- Stockwell, C.H., 1932, Great Slave Lake – Coppermine River area, Northwest Territories, *in* Geological Survey of Canada, Annual Report, Part C, p. 37-63.
- Stockwell, C.H., 1936, Eastern portion of Great Slave Lake, District of Mackenzie: Geological Survey of Canada, Maps 377A and 378A.
- Thomas, M.D., Gibb, R.A., and Quince, J.R., 1976, New evidence from offset aeromagnetic anomalies for transcurrent faulting associated with the Bathurst and McDonald faults, Northwest Territories: *Canadian Journal of Earth Sciences*, v. 13, p. 1244-1250, <http://dx.doi.org/10.1139/e76-126>
- Thompson, R.I., and Roots, C.F., 1982, Ogilvie Mountains Project, Yukon, *in* Current Research, Part A: Geological Survey of Canada, Paper 82-1A,

- p. 403-411.
- Thompson, R.I., Roots, C.F., and Mustard, P.S., 1994, Geology of Dawson map area (116B,C) northeast of Tintina Trench: Geological Survey of Canada, Open File 2849, <http://dx.doi.org/10.4095/194830>
- Umer, M., Kebede, S. and Osmaston, H., 2004, Quaternary glacial activity on the Ethiopian mountains, *in* Ehlers, J., and Gibbard, P.L., *eds.*, Quaternary Glaciations – Extent and Chronology, Part III: South America, Asia, Africa, Australasia, Antarctica: Elsevier, Amsterdam, p. 171-174, [http://dx.doi.org/10.1016/S1571-0866\(04\)80122-2](http://dx.doi.org/10.1016/S1571-0866(04)80122-2)
- Wallace, R.E., *ed.*, 1990, The San Andreas Fault System, California: United States Geological Survey, Professional Paper 1515, 283 p.
- Yeo, G.M., 1981, The Late Proterozoic Rapitan glaciation in the northern Cordillera, *in* Campbell, F.H.A., *ed.*, Proterozoic Basins of Canada: Geological Survey of Canada, Paper 81-10, p. 25-46.
- Zeller, S., 1987, Inventing Canada: Early Victorian Science and the Idea of a Transcontinental Nation: University of Toronto Press, Toronto, ON, 356 p.